

COVID-19 salivary signature: diagnostic and research opportunities

Dipak Sapkota,¹ Tine Merete Sølund,^{1,2} Hilde Kanli Galtung,¹ Lars Peter Sand,¹ Simone Giannechini,³ Kelvin K W To,^{4,5,6} Maria Cassia Mendes-Correa,⁷ Daniel Giglio,⁸ Bengt Hasséus,^{9,10} Paulo Henrique Braz-Silva ^{7,11}

For numbered affiliations see end of article.

Correspondence to

Professor Paulo Henrique Braz-Silva, Division of General Pathology, Department of Stomatology, Universidade de Sao Paulo Faculdade de Odontologia, Sao Paulo, Brazil; pbraz@usp.br and Professor Dipak Sapkota, Department of Oral Biology, Faculty of Dentistry, Oslo, Norway; dipak.sapkota@odont.uio.no

Received 3 June 2020
Revised 16 July 2020
Accepted 21 July 2020
Published Online First
7 August 2020

ABSTRACT

The COVID-19 (caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)) epidemic started in Wuhan (Hubei Province, China) in mid-December 2019 and quickly spread across the world as a pandemic. As a key to tracing the disease and to implement strategies aimed at breaking the chain of disease transmission, extensive testing for SARS-CoV-2 was suggested. Although nasopharyngeal/oropharyngeal swabs are the most commonly used biological samples for SARS-CoV-2 diagnosis, they have a number of limitations related to sample collection and healthcare personnel safety. In this context, saliva is emerging as a promising alternative to nasopharyngeal/oropharyngeal swabs for COVID-19 diagnosis and monitoring. Saliva collection, being a non-invasive approach with possibility for self-collection, circumvents to a great extent the limitations associated with the use of nasopharyngeal/oropharyngeal swabs. In addition, various salivary biomarkers including the salivary metabolomics offer a high promise to be useful for better understanding of COVID-19 and possibly in the identification of patients with various degrees of severity, including asymptomatic carriers. This review summarises the clinical and scientific basis for the potential use of saliva for COVID-19 diagnosis and disease monitoring. Additionally, we discuss saliva-based biomarkers and their potential clinical and research applications related to COVID-19.

INTRODUCTION

An epidemic of a new coronavirus with pneumonia-like symptoms started in Wuhan (Hubei Province, China) in December of 2019. The COVID-19, identified to be caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection,^{1,2} spread very quickly across the world and was declared a pandemic by WHO. As of 24 May 2020, the COVID-19 infection has accounted for >5 000 000 cases with >3 000 000 deaths reported worldwide.³ The fast spread of this disease is related to its highly infectious nature, and the disease is suggested to be transmitted through saliva droplets and nasal discharge.⁴ In order to trace the disease and to implement strategies aimed at breaking the chain of disease transmission, WHO has recommended extensive testing for COVID-19. This is particularly important as approximately 80% of the disease transmission has been reported to be related to asymptomatic cases.⁴ Here, we suggest that saliva-based testing can be an alternative to the

more widely used nasopharyngeal/oropharyngeal swabs for COVID-19 diagnosis and disease monitoring. In addition, we discuss unique opportunities and possible challenges related to the saliva-based research activities on COVID-19.

SALIVA AS A POTENTIAL DIAGNOSTIC FLUID FOR SARS-COV-2

The emergence of the COVID-19 pandemic has highlighted the need for multiple diagnostic strategies to efficiently evaluate potential cases in order to provide information on population exposure and immunity. These tools currently include virus molecular testing and rapid host immune response assays.

Saliva is a biological fluid in which SARS-CoV-2 can be found and for this reason saliva has been taken into consideration in the diagnosis of COVID-19. The presence of SARS-CoV-2 in saliva may be related to different sources such as i) virus entry to the oral cavity from lower/upper respiratory tract,^{1,5} ii) access to the mouth via oral cavity-specific crevicular fluid or iii) release of viral particles in the oral cavity via salivary ducts from the infected salivary glands⁶ (figures 1 and 2). The latter observation may explain how COVID-19 transmission can occur through asymptomatic cases with no obvious infection in the respiratory tracts.

The major salivary glands (parotid, submandibular and sublingual glands) are the major contributors of saliva secretion (figure 2). Approximately 600–1000 mL of saliva, containing molecules such as growth factors, cytokines and secretory IgA, is secreted each day from the human salivary glands.⁷ Of note, the unique salivary glands structure with rich surrounding blood circulation has been suggested to facilitate the exchange of molecules in the blood into the salivary acini and subsequently in the saliva.⁸ Saliva has been studied thoroughly as a potential diagnostic tool and it is expected to become a substitute for other biological fluids such as serum or urine in disease diagnosis.^{8,9} Compared with other diagnostic fluids, saliva sampling has the advantages and disadvantages as mentioned in table 1.

The diagnostic potential of saliva was established by studies that revealed that, like serum, saliva contains hormones, antibodies, growth factors, enzymes, microbes and their products that can enter saliva through blood via passive diffusion, active transport or extracellular ultra filtration. Therefore, saliva can be a reliable fluid for



© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Sapkota D, Sølund TM, Galtung HK, et al. *J Clin Pathol* 2021;**74**:344–349.




Figure 1 Schematic illustration demonstrating clinical implications and various sources of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in saliva. Inset: a suggested mechanism for SARS-CoV-2 entry into the salivary gland cells—the viral spike protein binds with the ACE2 receptor on the surface of the cell, followed by its priming with serine protease TMPRSS2 and subsequent entry into the cells. After replication and packaging, multiple new virus particles are released from the cells in saliva.

monitoring the physiological function of the body.¹⁰ Although the low concentration of some analytes in saliva compared with the blood previously proved challenging, the advent of highly sensitive molecular methods and nanotechnology have to a large extent circumvented this limitation.

Collection of saliva can be done in several ways, such as spitting out, collection with the help of sponge-like device and directly from the salivary gland duct.⁷ The spitting out technique is the cheapest one, and the saliva sample thus collected also includes nasopharyngeal/oropharyngeal/airway secretions. Sponge-like devices provide relatively more pure saliva, but this technique requires special equipment which is not always widely available. Saliva collected directly from the ducts of major salivary




Figure 2 Schematic illustration demonstrating major salivary glands (parotid, submandibular and sublingual) and their respective ducts, oropharynx and nasopharynx, and approximate anatomic locations for collection of oropharyngeal and nasopharyngeal swabs.

gland provides pure saliva, but the the collection process is time consuming and requires special equipment. Several protocols and approaches are available for DNA and RNA extraction and antibody detection, providing good performances regardless of sampling technique.¹¹

The diagnostic topic of saliva (called ‘Salivaomics’) includes the study of salivary proteins (proteomics), the study of salivary RNAs (transcriptomics), the study of salivary metabolites (metabolomics), the study of salivary microRNAs (microRNA) and the study of salivary microbiota (microbiome).¹²

To date, saliva is used for the diagnosis of several diseases including hereditary diseases, autoimmune diseases, malignancies, infections, dental caries and periodontal disease.^{13 14} Additionally, saliva can be used for diagnosing oral diseases with relevance for systemic diseases or for monitoring of levels of hormones, drugs and bone turnover markers.¹²

Diagnosis of saliva-based viral infections depends on the presence of viral DNA, RNA, microRNA, antigens or host antibodies in saliva. In this context, some viruses have been detected in saliva up to 29 days after infection, indicating that a saliva-based non-invasive diagnostic platform can be useful for early diagnosis and for monitoring the disease and treatment.^{9 15}

SALIVA TESTING FOR SARS-COV-2

SARS-CoV-2 detection using reverse-transcription PCR

SARS-CoV-2 is an enveloped, positive single-stranded RNA virus consisting of a core of RNA genome associated with nucleocapsid protein (N) and surrounded by a phospholipid membrane with three main viral structural proteins, spike surface glycoprotein (S), small envelope protein (E) and matrix protein (M).¹⁶ Nucleotide sequences within a number of SARS-CoV-2 genes such as E, RdRp, N1 and N2 and S can be used as detection targets for RT-PCR-based test methods.¹⁷ On the other hand, detection of SARS-CoV-2 antigens and/or immunoglobulins against them form the basis for enzyme immunoassays.¹⁷

Presently, RT-PCR is the most commonly used diagnostic test for the detection of SARS-CoV-2 RNA in the biological samples. For large-scale testing as in the case of SARS-CoV-2, proper selection of the type and the site of biological specimen collection is crucial for obtaining reliable test results.¹⁸ Biological samples from the upper (such as nasopharyngeal swabs, oropharyngeal swabs, throat swabs, nasal swabs) and lower (such as tracheal aspirates and bronchoalveolar lavage) respiratory tracts can be used for the detection of SARS-CoV-2 with varying degree of test sensitivity^{19–21} (figure 2). Tracheal aspirates and bronchoalveolar lavage, although more reliable for SARS-CoV-2 detection, are the less preferred specimens as compared with the nasopharyngeal/oropharyngeal swabs due to technical complexity in obtaining these samples.²¹ Currently, nasopharyngeal/oropharyngeal swabs where virus samples are collected by respectively rubbing the nasopharyngeal wall and the posterior pharynx/tonsillar areas with minitip swabs, are routinely used for SARS-CoV-2 detection¹⁷ (figure 2).

Despite the widespread use, the collection of nasopharyngeal/oropharyngeal swabs has a number of limitations.^{22 23} The collection of these swabs is less acceptable to patients as compared with non-invasive methods like saliva collection, as it tends to cause patient discomfort and even bleeding. Patient acceptance is highly desirable for test methods where multiple testing is needed for disease monitoring and follow-up, as in the case of COVID-19. Furthermore, the risk for disease transmission to the healthcare personnel when collecting these samples is high as it requires active involvement of the test taker. Additionally,

Table 1 Advantages and disadvantages of saliva sampling

Advantages	Disadvantages
Non-invasive approach for disease diagnosis and monitoring of general health. Painless (no patient discomfort and anxiety for sampling).	Not always reliable for measurement of certain markers. Contents of saliva can be influenced by the method of collection, degree of stimulation of salivary flow, interindividual variation and oral hygiene status.
Easy collection and applicable in remote areas. Relatively cheap technology.	Serum markers can reach whole saliva in an unpredictable way. Medications may affect salivary gland function and consequently the quantity and composition of saliva.
Cost-effective applicability for screening large populations. Suitable for children, anxious/disabled/elderly patients. Possible multisampling. Safer collection for health professionals than other biological samples such as nasopharyngeal swabs and blood. Cheap to store and ship. Easy to handle. No need for expensive equipment/instruments (swabs, suction tubes or special collection devices) for collection. Only needs a sterile container.	Possibility for degradation of salivary proteins due to presence of proteolytic enzymes.

collection of these samples demands the use of personal protective and healthcare resources, both of which tend to be in short supply in a pandemic like COVID-19.

Saliva as a biological fluid for molecular detection of SARS-CoV-2

Saliva is emerging as a promising alternative to nasopharyngeal/oropharyngeal swabs for COVID-19 diagnosis and monitoring.^{24 25} Indeed, the use of saliva as a biological specimen for SARS-CoV-2 testing to a great extent circumvents the above-mentioned limitations associated with the use of nasopharyngeal/oropharyngeal swabs. With clear instructions, patients can self-collect saliva samples. This is highly desirable in an outbreak in order to minimise the burden on healthcare personnel, the use of personal protective equipment and to allow serial sampling required for disease monitoring. A recent study has reported that self-collection of saliva sample for SARS-CoV-2 testing is feasible and can produce reliable test results.²⁴

The potential use of saliva for SARS-CoV-2 detection is scientifically well founded. Saliva is considered to be a good reservoir for viruses that originate from oral shedding, and secretions from the lower respiratory tract, nasopharynx and possibly infected salivary glands^{22 23} (figure 1). Indeed, Chen *et al* were able to detect SARS-CoV-2 RNA in three out of four saliva samples directly collected from the salivary gland ducts, thereby precluding contamination from respiratory secretions, of critically ill cases.²⁶ Together with the demonstration of ACE2 expression,²⁷ a main surface receptor type for SARS-CoV-2,¹ in the salivary gland, the above findings substantiate the idea that salivary gland could be one of the sources for SARS-CoV-2 in saliva. In line with this observation, recent studies by To *et al* demonstrated the presence of live SARS-CoV-2 in saliva.^{28 29} Furthermore, the possible diagnostic use of saliva for several respiratory viruses including coronavirus has been supported by studies demonstrating a high sensitivity and specificity of saliva-based tests, with >90% concordance between saliva and nasopharyngeal swabs.³⁰

Current studies from different groups have shown promising results on the possible use of saliva for detection of SARS-CoV-2 RNA^{24 25 28 29 31-33} (table 2). The sensitivity of saliva-based SARS-CoV-2 RNA detection methods seem to be comparable to^{24 32} or better than that of nasopharyngeal swabs.²⁵ Additionally, saliva seems to be a good candidate for the detection of

SARS-CoV-2 for cases with moderate-to-severe symptoms,²⁵ and for asymptomatic or mild cases.²⁴ The latter is particularly important for the screening of the suspicious/asymptomatic cases and for the surveillance of the healthcare workers. Self-sampling of saliva could also be an option in large-scale population-based point-prevalence studies.

Being a non-invasive specimen type, saliva is well-suited for serial viral load monitoring. The SARS-CoV-2 load in the saliva is reported to be highest after the first week of symptom onset, followed by a gradual decline.^{25 28 29} This underlines that saliva is a good candidate for SARS-CoV-2 detection in earlier disease phase. The temporal profile of SARS-CoV-2 load in saliva has been reported to be more consistent²⁵ as compared with that of nasopharyngeal swabs, suggesting its suitability for disease monitoring. Furthermore, saliva can be used for monitoring the response to antivirals in clinical trials.³⁴ Nonetheless, saliva can also be a potential source of viral transmission, thereby requiring standard protocols for its collection and subsequent handling. In the light of these promising results, a saliva-based SARS-CoV-2 RNA detection assay has already obtained approval through the US Food and Drug Administration emergency use authorisation.

Table 2 Main findings of recent studies on SARS-CoV-2 detection in saliva samples by using RT-PCR.

Authors	Main finding(s) related to salivary specimens
To <i>et al</i> ²⁸	91.7% of nasopharyngeal swab-diagnosed cases. Live virus was detected in saliva using viral culture.
To <i>et al</i> ²⁹	87% of nasopharyngeal swab-diagnosed cases. Salivary viral load was highest during the first week of symptom.
Azzi <i>et al</i> ³² Kojima <i>et al</i> ²⁴	Detected in all nasopharyngeal swab-diagnosed cases. Self-collected saliva and nasal swab had similar sensitivity as compared with the clinician-collected nasopharyngeal swabs
Wyllie <i>et al</i> ²⁵	Saliva is more sensitive and consistent than nasopharyngeal swabs
Williams <i>et al</i> ³¹	84.6% of nasopharyngeal swab-diagnosed cases. Viral load was higher in the nasopharyngeal swab.
Pasomub <i>et al</i> ³³	84.2% of nasopharyngeal swab-diagnosed cases. Saliva might be an alternative specimen for COVID-19 diagnosis.
SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.	

Antibodies against SARS-CoV-2

Besides RT-PCR-based RNA detection of SARS-CoV-2, preliminary studies have reported promising results for the detection of IgM and IgG against SARS-CoV-2 in serum/plasma samples of patients with SARS-CoV-2.³⁵ Interestingly, the production of SARS-CoV-specific secretory IgA in the saliva of mice intranasally immunised with SARS-CoV virus-like particles has been documented.³⁶ Hence, it is reasonable to speculate that anti-SARS-CoV-2 antibodies might also be present in human saliva.

Viral antibodies have been detected in saliva and the immunisation status of measles, rubella, mumps and hepatitis can be verified by analysing IgG, IgM and IgA in oral fluids.^{37 38} Regarding SARS-CoV-2, to date only a study protocol aimed to analyse IgG, IgM and IgA in different biological fluids including self-collected saliva for rapid SARS-CoV-2 diagnosis has been published.³⁹ However, there are so far no results describing the presence of antibodies against SARS-CoV-2 in human saliva. This clearly warrants future studies on the potential use of salivary immunoglobulins for COVID-19 in diagnostics, disease progression and immunisation monitoring.

Saliva biomarkers—perspectives for COVID-19 diagnosis and prognosis

Salivary biomarkers and their role in point-of-care applications have highlighted the development of the use of more advanced technologies such as micro/nanoelectro-mechanical systems, paper-based technology, RNA-sequencing, liquid biopsy, fluorescent biosensors, photometric and electrochemical methods, electric field-induced release and measurement method.^{12 40} Contemporary available point-of-care can be delivered in form of small and portable smartphones or 'lab-on-chips'.⁴⁰

Coronaviruses, such as SARS-CoV and Middle East respiratory syndrome (MERS)-CoV, have developed strategies to decrease or delay the production of interferon (IFN), triggering exuberant inflammatory responses leading to severe pulmonary conditions.^{41 42} The host's unregulated immune response and the production of inflammatory cytokines, known as 'cytokine storm', are believed to correlate with disease severity and poor prognosis during SARS-CoV and MERS-CoV infection.⁴¹ Several pro-inflammatory cytokines and chemokines, such as chemokine (C-C motif) ligand (CCL)-2, CCL-3, regulated on activation, normal T cell expressed and secreted (RANTES), interleukin (IL)-2 and IL-8, were highly expressed during MERS-CoV infection.⁴¹ Recent studies have reported that severe cases of COVID-19 exhibit increased plasma levels of IL-2, IL-6, IL-7, IL-10, granulocyte colony stimulating factor (G-CSF), INF- γ -inducible protein-10 (IP-10), macrophage chemotactic protein 1, macrophage inflammatory protein-1A and tumour necrosis factor- α compared with mild cases, indicating that the inflammatory response mediated by cytokine release is critical in the progression of COVID-19.^{42 43} Markers of the inflammatory process, such as cytokines and chemokines, can be measured in saliva. Such information has been suggested to be useful for the diagnosis and prognosis of both oral cavity and systemic diseases.¹³ Hence, it is possible to establish an inflammatory profile of COVID-19 by analysing inflammation-related biomarkers in saliva.

Unique proteomic, metabolic and/or lipid profiles in serum/plasma have been suggested to be useful in stratification of fatal/severe COVID-19 cases from the mild and healthy ones⁴⁴; and to predict progression of COVID-19 patients from a milder form to severe stages.⁴⁵ Interestingly, some of the identified biomarkers in these studies such as C reactive protein, lactate

dehydrogenase, malic acid, guanosine monophosphate and proteins associated with macrophage, platelet degranulation and complement system pathways are shown to be present in saliva. These findings support the possible use of saliva-based metabolic/protein/lipid biomarkers as a non-invasive approach for patient stratification in COVID-19 disease.

Metabolomics is a strategy used in the study of small molecules from the metabolic profile of cells, tissues or fluids, which help in the characterisation of a phenotype. These molecules, called biomarkers, are fundamental in clinical practice for determining the state of a disease.⁴⁶ Thus, metabolomics has helped to identify biomarkers with diagnostic potential and description of metabolic pathways in the most diverse clinical situations, including those involving viral and bacterial pathogens, and more specifically viruses that cause respiratory diseases such as influenza^{47 48} and SARS.⁴⁹ In a study by Wu *et al*,⁴⁹ patients recovered from severe acute respiratory syndrome caused by SARS-CoV were recruited after 12 years of infection for metabolic evaluation of the consequences of the disease. The comparison of patients' serum with healthy individuals showed differences in organic acids, amino acids, phospholipids, carnitine and inositol derivatives.⁴⁹ These results exemplify the practical application of metabolomics in the evaluation of long-term outcomes.

MicroRNAs, non-coding RNAs of 20-nucleotide to 22-nucleotide length, silencing gene expression by a transcript-specific target-mediate inhibitory activity, play a key role in several cellular processes including cell development and differentiation, immunity, cell metabolism, proliferation, apoptosis and cancer.⁵⁰ The relevance of monitoring microRNA is related to the fact that a single microRNA can be implicated in several cellular regulatory pathways, which involve different molecules. To date, there are studies reporting a particular microRNA upregulation and downregulation of nuclear factor- κ B pathway and IFN pathway associated with several viruses including respiratory virus infection.⁵¹ In this context, studies reporting coronavirus (including SARS-CoV) regulation of cellular microRNA showed the over-expression of miR-574-5p and miR214 and regulation of miR-9 and miR-98 with effect on apoptosis, cancer and autoimmune functions. Of note, SARS-CoV has been reported in the direct viral nucleocapsid downregulation of miR-223 and miR-98 expression, with effect in pro-inflammatory cytokine production.⁵² Additionally, in this context, since microRNAs associated with extracellular vesicles are known to be protected from enzymatic degradation, several studies have been focused on the investigation of the expression of microRNAs in extracellular vesicles obtained from saliva as potential biomarkers. Thus, the fact that microRNA present in biological fluid can reproduce the molecular event within the cellular context, make them a potential exhaustive marker to check the cell-infection status; this is especially important in a low replicative condition in which virus cannot be present in biological fluid,¹⁵ and provides an opportunity to assess virus pathological effect-associated diseases as in COVID-19.

Saliva mark of SARS-CoV-2 cell-receptor features in COVID-19

Its well known that SARS-CoV-2 infects host cells, including those in the respiratory tract lining, mainly using ACE2 receptor.⁵³ It is reported that SARS-CoV spike protein S has a high affinity for the ACE2 receptor and is activated by host type II transmembrane serine protease TMPRSS2 on primary target cells to fulfil viral entry⁵⁴ (figure 1). However, other host proteases such as furin on the tongue may be implicated in cleaving SARS-CoV-2

viral envelope glycoproteins in its furin-like cleavage site and enhancing infection with host cells.⁵⁵

The use of ACE2 receptor by other coronaviruses to infect salivary gland epithelial cells has been reported in rhesus macaques.⁶ Additionally, *in vitro* analysis of RNA-seq profiles from four public and consensus datasets revealed the expression of ACE2 receptor in human granular cells in salivary glands.²⁶ These observations suggest that the salivary glands can be a reservoir for SARS-CoV-2 and contribute to the presence of the transmissible form of viral infectious particles in saliva.⁵⁶ Additionally, it is possible that the salivary glands can harbour latent COVID-19 infection with possibility for subsequent reactivation. The latter suggestion clearly warrants further studies.

In addition to the salivary glands, ACE2 is abundantly expressed in the oral epithelial cells with highest expression in the tongue when compared with buccal and gingival tissues, T cells, B cells and oral fibroblasts.⁵⁷ These results raise a possibility that oral epithelial cells can function as a host for SARS-CoV-2. Epithelial cells in the oral mucosa are protected by a viscous mucous layer containing large glycoprotein macromolecules—mucins—produced in the salivary glands, and water. Virus particles must penetrate the mucous layer to be able to infect the cells in the epithelial lining. In the respiratory tract, mucins in the mucous layer have been shown to play a significant role in protecting airway epithelium to influenza and respiratory syncytial virus.⁵⁸ Surprisingly, few studies have looked at the role of mucous layer covering human epithelial linings during SARS-CoV-2 infection. Case reports have described mucous plugging in the lungs in postmortem examinations of patients that have succumbed to COVID-19.⁵⁹ Assuming that oral epithelial cells can be a possible route of entry for the SARS-CoV-2, studies are needed to understand the possible ways the virus particles penetrate the mucous layer and infect the underlying epithelial cells.

Of note, it has been reported by using airway epithelial cells *in vitro* that ACE2 is a human IFN-stimulated gene suggesting that SARS-CoV-2 could exploit IFN-driven upregulation of ACE2 as a mechanism to enhance viral infection and play a role in the development of COVID-19 pathogenesis.⁶⁰ All of these findings point out the potential interest in the investigation of saliva viral and host biomarkers as an opportunity to obtain a more complete molecular view of clinical relevance in the COVID-19 risk assessment as well as to develop new therapeutic antiviral treatments.

CONCLUSION AND FUTURE PERSPECTIVES

Saliva-based testing can be an alternative to the more widely used nasopharyngeal/oropharyngeal swabs for COVID-19 diagnosis and disease monitoring. The use of saliva-based SARS-CoV-2 testing offers several clinical advantages and is

scientifically well founded. However, further studies will be key to understanding the mutual relationship between COVID-19 and saliva, leading to the adoption of less invasive diagnostic techniques and facilitating the application of molecular tests on a large scale, a central strategy for controlling the epidemic. The search for salivary biomarkers associated with the development and progression of COVID-19 could allow a better distinction between asymptomatic, mild, moderate or advanced disease. Knowledge of this kind might lead to the development of point-of-care devices, which can be extremely useful for understanding of the evolution of contagions and immunological responses in population studies.

Author affiliations

- ¹Department of Oral Biology, Faculty of Dentistry, University of Oslo, Oslo, Norway
- ²Department of Pathology, Rikshospitalet University Hospital, Oslo, Norway
- ³Department of Experimental and Clinical Medicine, Università degli Studi di Firenze, Firenze, Toscana, Italy
- ⁴State Key Laboratory for Emerging Infectious Diseases, Department of Microbiology, Li KaShing Faculty of Medicine of the University of Hong Kong, Hong Kong, China
- ⁵Department of Microbiology, Queen Mary Hospital, Hong Kong, China
- ⁶Department of Clinical Microbiology and Infection Control, The University of Hong Kong – Shenzhen Hospital, Shenzhen, China
- ⁷Laboratory of Virology (LIM-52), Institute of Tropical Medicine of São Paulo, School of Medicine, University of São Paulo, São Paulo, Brazil
- ⁸Department of Clinical Oncology, The Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden
- ⁹Department of Oral Medicine and Pathology, Institute of Odontology, The Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden
- ¹⁰Clinic of Oral Medicine, Region Västra Götaland, Gothenburg, Sweden
- ¹¹Department of Stomatology, School of Dentistry, University of São Paulo, São Paulo, Brazil

Handling editor Tahir S Pillay.

Acknowledgements The authors would like to thank Evan Michael Vallenari for language correction of the manuscript.

Contributors DS, SG, BH and PHB-S contributed to conception, design, data interpretation and drafted the manuscript; TMS, HKG, LPS, KKWT, MCM-C and DG contributed to design and data interpretation, critically revised the manuscript. All authors gave their final approval and agreed to be accountable for all aspects of the work.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iD

Paulo Henrique Braz-Silva <http://orcid.org/0000-0002-1842-9521>

REFERENCES

- 1 Zhou P, Yang X-L, Wang X-G, *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;579:270–3.
- 2 Coronaviridae Study Group of the International Committee on Taxonomy of Viruses. The species severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol* 2020;5:536–44.
- 3 WHO. World Health organization. coronavirus disease (COVID-2019) situation report-125. secondary World Health organization. coronavirus disease (COVID-2019) situation report-125, 2020. Available: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/>
- 4 Li R, Pei S, Chen B, *et al.* Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). *Science* 2020;368:489–93.
- 5 Zhu N, Zhang D, Wang W, *et al.* A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med Overseas Ed* 2020;382:727–33.
- 6 Liu L, Wei Q, Alvarez X, *et al.* Epithelial cells lining salivary gland ducts are early target cells of severe acute respiratory syndrome coronavirus infection in the upper respiratory tracts of rhesus macaques. *J Virol* 2011;85:4025–30.
- 7 Malamud D. Saliva as a diagnostic fluid. *Dent Clin North Am* 2011;55:159–78.
- 8 Zhang C-Z, Cheng X-Q, Li J-Y, *et al.* Saliva in the diagnosis of diseases. *Int J Oral Sci* 2016;8:133–7.

Take home messages

- ▶ Saliva has been described as a good candidate for diagnosis of COVID-19, showing sensitivity and specificity comparable to nasopharyngeal swabs.
- ▶ Saliva is a non-invasive, easy to handle and with the possibility of self-collection body fluid—these characteristics are important in a pandemic scenario, leading to less exposure of healthcare professionals.
- ▶ Saliva biomarkers have a potential to be an important guide in COVID-19 prognosis, making possible the development of point-of-care devices.

- 9 Castro T, Sabalza M, Barber C, et al. Rapid diagnosis of Zika virus through saliva and urine by loop-mediated isothermal amplification (lamp). *J Oral Microbiol* 2018;10:1510712.
- 10 Lima DP, Diniz DG, Moimaz SAS, et al. Saliva: reflection of the body. *Int J Infect Dis* 2010;14:e184–8.
- 11 Portilho MM, Mendonça ACF, Marques VA, et al. Comparison of oral fluid collection methods for the molecular detection of hepatitis B virus. *Oral Dis* 2017;23:1072–9.
- 12 Kaczor-Urbanowicz KE, Martin Carreras-Presas C, Aro K, et al. Saliva diagnostics - Current views and directions. *Exp Biol Med* 2017;242:459–72.
- 13 Galhardo LF, Ruivo GF, de Oliveira LD, et al. Inflammatory markers in saliva for diagnosis of sepsis of hospitalized patients. *Eur J Clin Invest* 2020;50:e13219.
- 14 Marques Filho JS, Gobara J, da Silva Salomao GV, et al. Cytokine levels and human herpesviruses in saliva from clinical periodontal healthy subjects with peri-implantitis: a case-control study. *Mediators Inflamm* 2018;2018:1–7.
- 15 Martelli F, Mencarini J, Rocca A, et al. Polyomavirus microRNA in saliva reveals persistent infectious status in the oral cavity. *Virus Res* 2018;249:1–7.
- 16 Wu C, Liu Y, Yang Y, et al. Analysis of therapeutic targets for SARS-CoV-2 and discovery of potential drugs by computational methods. *Acta Pharm Sin B* 2020;10:766–88.
- 17 Pang J, Wang MX, Ang IYH, et al. Potential rapid diagnostics, vaccine and therapeutics for 2019 novel coronavirus (2019-nCoV): a systematic review. *J Clin Med* 2020;9:623.
- 18 Xie C, Jiang L, Huang G, et al. Comparison of different samples for 2019 novel coronavirus detection by nucleic acid amplification tests. *Int J Infect Dis* 2020;93:264–7.
- 19 Lin C, Xiang J, Yan M, et al. Comparison of throat swabs and sputum specimens for viral nucleic acid detection in 52 cases of novel coronavirus (SARS-CoV-2)-infected pneumonia (COVID-19). *Clin Chem Lab Med* 2020;58:1089–94.
- 20 Wu J, Liu J, Li S, et al. Detection and analysis of nucleic acid in various biological samples of COVID-19 patients. *Travel Med Infect Dis* 2020:101673.
- 21 Liu R, Han H, Liu F, et al. Positive rate of RT-PCR detection of SARS-CoV-2 infection in 4880 cases from one hospital in Wuhan, China, from Jan to Feb 2020. *Clin Chim Acta* 2020;505:172–5.
- 22 Henrique Braz-Silva P, Pallos D, Giannecchini S, et al. SARS-CoV-2: what can saliva tell us? *Oral Dis* 2020. doi:10.1111/odi.13365. [Epub ahead of print: 20 Apr 2020].
- 23 Sapkota D, Thapa SB, Hasséus B, et al. Saliva testing for COVID-19? *Br Dent J* 2020;228:658–9.
- 24 Kojima NTF, Slepnev V, Bacelar A, et al. Self-Collected oral fluid and nasal swabs demonstrate comparable sensitivity to clinician collected nasopharyngeal swabs for Covid-19 detection. *medRxiv* 2020. [Epub ahead of print: 15 Apr 2020].
- 25 Wyllie AL FJ, Casanovas-Massana A, Campbell M, et al. Saliva is more sensitive for SARS-CoV-2 detection in SARS-CoV-2 patients than nasopharyngeal swabs. *medRxiv* 2020. [Epub ahead of print: 22 Apr 2020].
- 26 Chen LZJ, Peng J, Li X, et al. Detection of 2019-nCoV in saliva and characterization of oral symptoms in COVID-19 patients. *SSRN* 2020.
- 27 Chen Y, Liu Q, Guo D. Emerging coronaviruses: genome structure, replication, and pathogenesis. *J Med Virol* 2020;92:418–23.
- 28 To KKW, Tsang OT-Y, Leung W-S, et al. Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study. *Lancet Infect Dis* 2020;20:565–74.
- 29 To KK-W, Tsang OT-Y, Chik-Yan Yip C, et al. Consistent detection of 2019 novel coronavirus in saliva. *Clin Infect Dis* 2020. doi:10.1093/cid/ciaa149. [Epub ahead of print: 12 Feb 2020].
- 30 To KKW, Yip CCY, Lai CYW, et al. Saliva as a diagnostic specimen for testing respiratory virus by a point-of-care molecular assay: a diagnostic validity study. *Clin Microbiol Infect* 2019;25:372–8.
- 31 Williams E, Bond K, Zhang B, et al. Saliva as a noninvasive specimen for detection of SARS-CoV-2. *J Clin Microbiol* 2020;58.
- 32 Azzi L, Carcano G, Gianfagna F, et al. Saliva is a reliable tool to detect SARS-CoV-2. *J Infect* 2020;81:e45–50.
- 33 Pasomsub E, Watcharananan SP, Boonyawat K, et al. Saliva sample as a non-invasive specimen for the diagnosis of coronavirus disease 2019: a cross-sectional study. *Clin Microbiol Infect* 2020. doi:10.1016/j.cmi.2020.05.001. [Epub ahead of print: 15 May 2020].
- 34 Hung IF-N, Lung K-C, Tso EY-K, et al. Triple combination of interferon beta-1b, lopinavir-ritonavir, and ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label, randomised, phase 2 trial. *Lancet* 2020;395:1695–704.
- 35 Zhao J, Yuan Q, Wang H, et al. Antibody responses to SARS-CoV-2 in patients of novel coronavirus disease 2019. *Clin Infect Dis* 2020. doi:10.1093/cid/ciaa344. [Epub ahead of print: 28 Mar 2020].
- 36 Lu B, Huang Y, Huang L, et al. Effect of mucosal and systemic immunization with virus-like particles of severe acute respiratory syndrome coronavirus in mice. *Immunology* 2010;130:254–61.
- 37 Cruz HM, de Paula VS, da Silva EF, et al. Utility of oral fluid samples for hepatitis B antibody detection in real life conditions. *BMC Infect Dis* 2019;19:632.
- 38 Sampaio BCF, Rodrigues JP, Meireles LR, et al. Measles, rubella, mumps and Toxoplasma gondii antibodies in saliva of vaccinated students of schools and universities in São Paulo City, Brazil. *Braz J Infect Dis* 2020;24:51–7.
- 39 Sullivan PS, Sailey C, Guest JL, et al. Detection of SARS-CoV-2 RNA and antibodies in diverse samples: protocol to validate the sufficiency of Provider-Observed, Home-Collected blood, saliva, and oropharyngeal samples. *JMIR Public Health Surveill* 2020;6:e19054.
- 40 Aro K, Wei F, Wong DT, et al. Saliva liquid biopsy for point-of-care applications. *Front Public Health* 2017;5:77.
- 41 Fehr AR, Channappanavar R, Perlman S. Middle East respiratory syndrome: emergence of a pathogenic human coronavirus. *Annu Rev Med* 2017;68:387–99.
- 42 Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;395:507–13.
- 43 Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497–506.
- 44 Wu D, Shu T, Yang X, et al. Plasma metabolomic and lipidomic alterations associated with COVID-19. *Nat Sci Rev* 2020. [Epub ahead of print: 28 Apr 2020].
- 45 Shen B, Yi X, Sun Y, et al. Proteomic and metabolomic characterization of COVID-19 patient sera. *Cell* 2020;182:59–72.
- 46 Goldansaz SA, Guo AC, Sajed T, et al. Livestock metabolomics and the livestock metabolome: a systematic review. *PLoS One* 2017;12:e0177675.
- 47 Chandler JD, Hu X, Ko E-J, et al. Metabolic pathways of lung inflammation revealed by high-resolution metabolomics (HRM) of H1N1 influenza virus infection in mice. *Am J Physiol Regul Integr Comp Physiol* 2016;311:R906–16.
- 48 Banoei MM, Vogel HJ, Weljje AM, et al. Plasma metabolomics for the diagnosis and prognosis of H1N1 influenza pneumonia. *Crit Care* 2017;21:97.
- 49 Wu Q, Zhou L, Sun X, et al. Altered lipid metabolism in recovered SARS patients twelve years after infection. *Sci Rep* 2017;7:9110.
- 50 O'Brien J, Hayder H, Zayed Y, et al. Overview of microRNA biogenesis, mechanisms of actions, and circulation. *Front Endocrinol* 2018;9:402.
- 51 Leon-Icaza SA, Zeng M, Rosas-Taraco AG. microRNAs in viral acute respiratory infections: immune regulation, biomarkers, therapy, and vaccines. *ExRNA* 2019;1.
- 52 Canatan D, De Sanctis V. The impact of microRNAs (miRNAs) on the genotype of coronaviruses. *Acta Biomed* 2020;91:195–8.
- 53 Wrapp D, Wang N, Corbett KS, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 2020;367:1260–3.
- 54 Song J, Li Y, Huang X, et al. Systematic analysis of ACE2 and TMPRSS2 expression in salivary glands reveals underlying transmission mechanism caused by SARS-CoV-2. *J Med Virol* 2020. [Epub ahead of print: 20 May 2020].
- 55 Coutard B, Valle C, de Lamballerie X, et al. The spike glycoprotein of the new coronavirus 2019-nCoV contains a furin-like cleavage site absent in CoV of the same clade. *Antiviral Res* 2020;176:104742.
- 56 Xu J, Li Y, Gan F, et al. Salivary glands: potential reservoirs for COVID-19 asymptomatic infection. *J Dent Res* 2020;99:989.
- 57 Xu H, Zhong L, Deng J, et al. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int J Oral Sci* 2020;12:8.
- 58 Zanin M, Baviskar P, Webster R, et al. The interaction between respiratory pathogens and mucus. *Cell Host Microbe* 2016;19:159–68.
- 59 Barton LM, Duval EJ, Stroberg E, et al. COVID-19 autopsies, Oklahoma, USA. *Am J Clin Pathol* 2020;153:725–33.
- 60 Ziegler CGK, Allon SJ, Nyquist SK, et al. SARS-CoV-2 receptor ACE2 is an interferon-stimulated gene in human airway epithelial cells and is detected in specific cell subsets across tissues. *Cell* 2020;181:1016–35.