

Supplemental Table 1: Summary of established and novel formulae to calculate LDL-cholesterol

Author (ref)	Year	Setting	N	Exclusions	Patients	Fasting status	Limitations	Formula	Performance
<i>Friedewald</i> <sup>1</sup>	1972	United States	448	TG>4.5 mmol/L Non-fasted patients	Familial Hyper-Cholesterolaemia	Fasted 12-14h	TG>4.5mmol/L LDL<1.8mmol/L	LDL-C = TC – HDL-C – TG/2.2 (mmol/L)	High correlation coefficients ranging from 0.94 to 0.98 (healthy population) and 0.99 (patients with dyslipidaemia)
<i>DeLong</i> <sup>2</sup>	1986	North America	>10000	Not available	Not available	Fasted ≥ 12hrs	Not available	LDL-C = TC – (HDL-C + 0.37 × TG)	Not available
<i>Rao</i> <sup>3</sup>	1988	Kuwait	196	None	Serial Lipid profiles	Unknown	None	$\frac{[(\text{HDL-C} \times (a - c)] + (\text{TC} \times c) - \text{TG}}{(c - b)}$ <p>A= HDL: TG/TC ratio = 0.15 B= LDL: TG/TC ratio = 0.095 C= VLDL: TG/TC ratio = 2.1</p> <p>Ratios are determined using results from ultracentrifugation fractions</p>	Linear correlation coefficients ranged between 0.93 – 0.937

<i>Hattori</i> <sup>4</sup>	1998	Japan	2179	TC<2.5 and >9 mmol/L and and TG<0.3 and >4.5 mmol/L Apo B <0.7umol/L	2161 out- and inpatients and 18 healthy controls	Fasted overnight	TC>9mmol/L and hyper-triglyceridaemia	LDL-C = (0.94TC – 0.94HDL-C – 0.19TG)	( <i>r</i> = 0.86) Between estimated and measured LDL was observed.
<i>Puavilai</i> <sup>5</sup>	2004	Thailand	1079	NA	Serum samples	Fasted	NA	LDL-C = TC – HDL-C – (TG/6)	Friedewald (72.3%) vs Puavillai (91.6%)
<i>Anandaraja</i> <sup>6</sup>	2005	India	2008	Not stated	Blood samples-lipid analysis (Indian patients)	Unknown	None	LDL-C = (0.9TC) – (0.9TG/5) – 28	d-LDL-C, LDL-C shows good correlation( <i>r</i> =0.97)
<i>Teerakanchana</i> <sup>7</sup>	2007	Thailand	1016	None	Outpatients Aged 8-89	Fasted 12-14 hrs	None	LDL-C = 0.91TC – 0.634(HDL-C) – 0.111TG – 6.755	Max correlation coefficient <i>r</i> =980

<i>Ahmadi</i> <sup>8</sup>	2008	Iran	237	None	TC>6.46mmol/L	Fasted	None	$LDL-C = (TC/1.19) - (HDL-C/1.1) - (TG/1.9) - 38$	No significant difference between direct and Ahmadi methods.
<i>Chen</i> <sup>9</sup>	2010	China	2180	<4.5 mmol/L	Outpatients >18 years	Fasted overnight	<4.5mmol/L	$LDL-C = 0.9(Non-HDL-C) - 0.1TG$ (mg/dL)	Improved vs FW
<i>Vujovic</i> <sup>10</sup>	2010	Serbia	2053	TG>4.5 mmol/L	Routine follow up patients	Fasted overnight	TG >4.5 mmol/L	$LDL-C = TC - TG/3 - HDL-C$ (mmol/L)	No significant difference between Vujovic LDL-C and direct-LDL was found.
<i>De Cordova</i> <sup>11</sup>	2013	Brazil	10664	None	Patients between 1-93 years old	Fasted 12 hrs	Not stated	$LDL-C = 0.75(TC - HDL-C)$	Higher correlation with directly measured LDL compared with Friedewald

<i>Martin/Hopkins</i> <sup>12</sup>	2013	United States	1350908	None	Children adolescents and adults	Unknown	TG < 4.5 mmol/L	LDL-C = TC – HDL-C – TG/adjustable factor	MH > FW for TG > 4 mmol/L and LDL < 1.8 mmol/L vs dLDL
<i>Sampson</i> <sup>13</sup>	2020	United States	8656	Results < 1st and > 99th percentiles for each lipid parameter	Fasting patients at the NIH clinical centre	Unknown	TG > 9 mmol/L	LDL-C (mg/dL) = TC/0.948 – HDL-C/0.971 – (TG/8.56 + TG × non-HDL-C/2140 – TG <sup>2</sup> /16100) – 9.44	35% less misclassifications in patients with hypertriglyceridemia.
<i>Dudi</i> <sup>14</sup>	2022	Sub-Himalayan region	1851	TG > 400 mg/dL / 4.5 mmol/L	Fasting lipid profile in a Sub-himalayan population	Unknown	Does not account for general population and dylipidaemia	LDL-C = -2.421 + (0.752 × TC) – (0.047 × TG) – (0.350 × HDL-C)	Showed a better diagnostic performance as compared to other 10 equations taking Direct LDL-C as gold standard for Sub-himalayan Population and may be used as a substitute for FW in the study population.
<i>Sampson</i> <sup>15</sup>	2022		24713	TG > 3000 mg/dL / 33.87 mmol/L (N = 38) TG > 1500 mg/dL / 16.9 mmol/L (N = 3) Samples with detectable	Dyslipidaemic patients		An additional apoB test is required Equation only compared the original Sampson equation and eS-VLDL-C to F-LDL-C and did	eS-VLDL-C = (non-HDL-C/3.81) – (HDL-C/8.93) + (TG/7.73) + [(non-HDL-C × TG)/2050] – (TG <sup>2</sup> /13300) – (ApoB/2.49) – [(ApoB × TG)/3550] + 7.46	At a cut-point of 0.194, the first method showed a modest ability for identifying HLP3 (sensitivity = 73.9%; specificity = 82.6%; and AUC = 0.8685) but was comparable to the existing dual lipid apoB ratio method.

		United States		lipoprotein – X on agarose gel electrophoresis (N = 126)		Unknown	not include other equations  New equation not assessed in the routine clinical lab setting  No variability in ethnic groups		The second method based on eS-VLDL-C showed much better sensitivity (96.5%) and specificity (94.5%) at a cut-point of 0.209. It also had an excellent AUC value of 0.9912 and was superior to the two other methods in test classification.
--	--	---------------	--	--	--	---------	---	--	---

Supplemental Table 2: Comparison of novel LDL-C equations in different populations

Country/Region	Clinical interest group	Homogeneous assay used	Calculations used	Publication year	Conclusion
<b>Brazil</b>	Type II Diabetes	Unspecified	Friedewald, Martin/Hopkins, Sampson/NIH	2022	Martin/Hopkins and Sampson's equations show a similar accuracy for calculating LDL-C concentrations in patients with diabetes, and both equations were marginally better than the Friedewald equation <sup>16</sup>
<b>China</b>	Coronary artery disease	Selective solubilization method test kit: <i>Kyowa Medex</i>	Friedewald, Martin/Hopkins, Sampson/NIH	2022	Even though misclassification by the Martin/Hopkins and Sampson equations was present in approximately 20% of patients versus directly measured LDL-C, directly measuring LDL-C did not identify any more individuals with increased risk of all-cause mortality, recurrent myocardial infarction, and major adverse cardiovascular events <sup>17</sup>
<b>Germany</b>	Chronic kidney disease	LDL-Cholesterol Gen. 3 kit on the Cobas VR modular platform 8000 or 6000: <i>Roche Diagnostics</i>	Friedewald, Martin/Hopkins	2020	The Friedewald formula yields lower LDL-C concentrations in CKD than direct enzymatic measurements <sup>18</sup>
<b>Ghana</b>	Cardiothoracic disease	Chema LDL direct FL test: <i>Hospitex Diagnostics</i>	Friedewald, Anandaraja, Martin-Hopkins	2018	The modified LDL-C formula derived by this study was more accurate compared to Anandaraja and Friedewald and could serve as a substitute for direct LDL-C and Friedewald in this population <sup>19</sup>
<b>Hungary</b>	All patients	Direct homogeneous method Wako on AU5800: <i>Beckman-Coulter</i>	Friedewald, Martin/Hopkins	2020	LDL-C estimation using the Martin/Hopkins formula yields a more accurate LDL-C value than that calculated by the Friedewald formula in this population <sup>20</sup>

<b>India</b>	All patients	LDL-C plus 2nd generation assay: <i>Roche Diagnostics</i>	Friedewald, Martin-Hopkins	2021	Martin/Hopkins calculated LDL-C values showed better agreement with directly measured values, and less bias and percentage total error than those obtained by use of the Friedewald formula <sup>21</sup>
	Hyperlipidaemia	Enzymatic colorimetric assay with rapid reagent kit: <i>Agappe</i>	Friedewald	2021	Both direct and calculated methods of LDL estimation have their limitations <sup>22</sup>
	Type II Diabetes	AU 680 clinical chemistry analyzer: <i>Beckmann-Coulter</i>	Friedewald, Hattori, Anandaraja, Chen, de Cordova, Teerakanchana, Ahmadi, DeLong, Rao, Martin/Hopkins, Puavilai, Hata, Vujovic	2022	In this population all formulae except Ahmadi significantly underestimated the LDL-C when compared with the direct assay. The newer Martin's formula appeared to more precisely calculate LDL-C when compared with the Friedewald formula <sup>23</sup>
<b>Iran</b>	Adults	Direct LDL-C kit from Pars Azmoon on Roche Hitachi 912 chemistry analyser: <i>Roche Diagnostics</i>	Friedewald, Hattori, Anandaraja, Chen, de Cordova, Teerakanchana, Ahmadi, DeLong, Rao, Puavilai Vujovic	2020	The Friedewald formula can be used in serum samples with TG up to 400 mg/dL when validated methods with correct calibration and control systems are used to derive the formula's componetns <sup>24</sup>
	Children	Enzymatic method on Auto Analyzer: <i>Hitachi</i>	Friedewald, Ahmadi, Chen, Anandaraja	2020	Friedewald had the strongest correlation with the direct assay and the Ahmadi formula the weakest at TG cut-ff of 100 mg/dL <sup>25</sup>
<b>Israel</b>	PCSK-9 inhibitors	N/A	Friedewald, Martin/Hopkins	2021	In high-risk patients achieving low LDL-C on PCSK9i treatment, Martin/Hopkins displayed significant proportion of LDL-C upward discordance compared to the Friedewald

					equation, identifying patients that may need treatment intensification <sup>26</sup>
	Coronary artery disease	N/A	Friedewald, Martin/Hopkins, Sampson/NIH	2020	Both Martin/Hopkins and Sampson/NIH LDL-C estimates displayed significant proportion of upward discordance compared to Friedewald formula, particularly in patients with elevated triglycerides and low LDL-C <sup>27</sup>
<b>Italy</b>	Children	Colorimetric Enzymatic Selective elimination assay: <i>Beckman-Coulter</i>	Friedewald; Sampson/NIH	2021	Sampson/NIH equation is more reliable than Friedewald <sup>28</sup>
	Hypertension	N/A	Friedewald, Martin/Hopkins, Sampson/NIH	2021	Higher LDL-C control rates were achieved using the Martin/Hopkins formula <sup>29</sup>
<b>Japan</b>	Type II Diabetes, Hypertriglyceridaemia	Unspecified	N/A	2022	The TG/LDL-C ratio is a reliable surrogate lipid marker of small dense LDL-C and superior to non-HDL-C in type 2 diabetes patients not treated with statins <sup>30</sup>
<b>Korea</b>	All patients	Pureauto S Cholestest-LDL on Automatic Analyzer 7600: <i>Hitachi</i>	Friedewald	2020	The Friedewald equation tends to underestimate LDL-C in high-risk subjects such as hypertriglyceridemia and hypo-HDL-cholesterolemia <sup>31</sup>
	Adults	Reagents from Sekisui, Beckman, or Roche Diagnostics on Hitachi 7600, Modular D2400, Beckman AU5800 or Abbott Architect Ci8200 automated analysers	Friedewald, Martin/Hopkins, Sampson/NIH	2021	Martin/Hopkins may be a readily adoptable and cost-effective alternative to direct LDL-C measurement, irrespective of dyslipidaemia status <sup>32</sup>



	Adults	LDL-cholesterol plus 2nd generation reagent and LDL-cholesterol Gen.3 on Cobas 8000 c702: Roche Diagnostics	Friedewald, Hattori, Anandaraja, Chen, De Cordova, Ahmadi, Delong, Rao, Puavilai, Martin/Hopkins, Vujovic	2021	The accuracy of 12 equations for LDL-C varied by cohort and subgroup based on direct-LDL and triglycerides <sup>33</sup>
	Atherosclerosis	Unspecified	Friedewald, Martin/Hopkins	2018	The Martin/Hopkins equation showed a 2-fold better concordance with LDL-C measured with a direct homogeneous assay in coronary atherosclerosis compared to the Friedewald equation when the TG level was $\geq 200$ mg/dL <sup>34</sup>
<b>Lebanon</b>	All patients	Colorimetric assay after precipitation of non-LDL lipoproteins on 5600 dry chemistry analyzer: VITROS	Friedewald, Martin/Hopkins, de Cordova, Sampson/NIH	2022	A machine learning algorithm using K nearest neighbours is promising for the estimation of LDL-C as it agrees better with directly measured LDL-C than equations, especially in mild and severe hypertriglyceridemia <sup>35</sup>
<b>Nigeria</b>	Adults	Unspecified	Friedewald, Martin/Hopkins	2021	The Friedewald equation results in a lower LDL-C than the Martin/Hopkins formula <sup>36</sup>
<b>North America</b>	Adults	N/A	Friedewald, Martin/Hopkins	2021	The Martin/Hopkins equation correlates more strongly with non-HDL-C and ApoB than Friedewald and Friedewald underestimates LDL-C at low concentrations <sup>37</sup>
	PCSK-9 inhibitors	N/A (beta-quantification used)	Friedewald, Martin-Hopkins, Sampson/NIH	2021	LDL-C calculated by any of the three methods can guide treatment decisions for most patients, including those treated with proprotein convertase subtilisin/kexin type 9 inhibitors <sup>38</sup>
	Hypertriglyceridaemia	N/A (Vertical Auto Profile ultracentrifugation used)	Friedewald, Martin/Hopkins, Extended Martin/Hopkins, Sampson/NIH	2021	The extended Martin/Hopkins equation offered greater LDL-C accuracy compared with the Friedewald and Sampson/NIH equations in patients with triglyceride levels of 400 to 799 mg/dL <sup>39</sup>

Adults	N/A	Friedewald, Martin/Hopkins, Sampson/NIH	2021	The Martin/Hopkins Sampson/NIH equations did not over- or under-estimate ASCVD risk compared to Friedewald in primary prevention, while Friedewald underestimated subclinical ASCVD risk in the low-risk population. Martin/Hopkins or Sampson/NIH should be used for lipid screening in the general population <sup>40</sup>
Type II Diabetes > 40 yrs	N/A	Friedewald, Martin/Hopkins	2020	The Martin/Hopkins equation for estimating LDL-C is a significant improvement on the Friedewald formula but remains an imperfect tool to estimate the atherogenic load in patients with high TG levels. Concordance with the alternate target parameters (apoB and non-HDL-C) was higher for Martin/Hopkins than for Friedewald when triglycerides levels were increased <sup>41</sup>
Hypertriglyceridaemia in adults	Enzymatic colorimetric assay on 8000 Chemistry analyzer: <i>Roche Diagnostics</i>	Friedewald; Martin/Hopkins, Sampson	2022	Martin/Hopkins reduces LDL-C treatment group miscategorization rate leading to fewer underestimations of risk overall compared to Sampson/NIH; however, neither may be sufficiently accurate to report LDL-C in patients with TG $\geq 4.52$ mmol/L ( $\geq 400$ mg/dL) <sup>42</sup>
Females with HIV	2-phase enzymatic assay after selectively solubilizing and enzymatically digesting non-LDL lipoproteins: <i>Quest diagnostics</i>	Friedewald, Martin/Hopkins, Sampson/NIH	2021	All 3 calculation methods have lower accuracy in women with HIV, even after adjusting for TG levels and fasting status <sup>43</sup>
Males with HIV	N/A	Friedewald, Martin/Hopkins	2020	The Martin/Hopkins equation is more accurate than the Friedewald equation, especially at higher triglyceride levels, which are more common in people with HIV <sup>44</sup>

	Dyslipidaemia	N/A (beta-quantification used)	Modified Sampson/NIH	2022	The Sampson/NIH equation modified to include apoB, is nearly as accurate as the beta quantification reference method for the diagnosis of hyperlipoproteinaemia type 3 <sup>15</sup>
<b>Slovakia</b>	All patients	N/A	Friedewald, Martin/Hopkins, Sampson/NIH	2021	The results of clinical and epidemiological studies are significantly influenced by the method used to determine LDL-C <sup>45</sup>
<b>South Africa</b>	Outpatients	Automated direct enzymatic LDL-C: <i>Abbott Architect, Roche Cobas</i>	Friedewald, Hattori, Anandaraja, Chen, De Cordova, Sampson/NIH, Ahmadi, Delong, Martin/Hopkins, Puavilai, Vujovic	2021	Replacement of the Friedewald equation with Martin/Hopkins estimation to improve quality of LDL-C results can be safely implemented across analysers <sup>46</sup>
	Children Adults with diabetes	Automated direct enzymatic LDL-C: <i>Abbott Architect, Roche Cobas</i>	Friedewald, Hattori, Anandaraja, Sampson/NIH, Martin/Hopkins, Extended Martin/Hopkins	2022	Different measurement platforms influence the results of predictive equations and directly measured LDL-C: propose utilizing the Martin-Hopkins equation as an alternative to dLDL-C assays in adults with diabetes and for screening purposes in paediatric populations to avoid underestimating cardiovascular risk [94]
<b>Sub-Himalayan</b>	All patients	Direct LDL-C on AU5800 Clinical Chemistry Automated System: <i>Beckman-Coulter</i>	Friedewald, Puavilai, Vujovic, Anandaraja, Gowda, Teerakanchana, Ahmadi, de Cordova, Chen, Hattori	2022	The equation derived in this study showed a better diagnostic performance as compared to 10 other equations taking Direct LDL-C as the gold standard for Sub-Himalayan Population and may be used as a substitute for Friedewald <sup>14</sup>
<b>Thailand</b>	Adults	Unspecified	Friedewald, Ahmadi,	2021	Friedewald, Sampson/NIH, and Chen equations showed better estimations of LDL-C levels and

			Anandaraja, Chen, Cordova, Densethakul, Hattori, Martin, Puavillai, Sampson, Vujovic		had higher accuracy in classifying the degree of LDL-C strata <sup>47</sup>
<b>Turkey</b>	Cardiovascular disease > 40 yrs	N/A	Friedewald, Martin/Hopkins, Sampson/NIH	2021	In most cardiology outpatients, the Friedewald equation has excellent agreement with the Martin/Hopkins and Sampson equations <sup>48</sup>
	Adults with normotriglyceridaemia	2nd generation reagent on the 6000 c501 analyzer: <i>Roche Diagnostics</i>	Friedewald, Martin/Hopkins	2021	The Martin/Hopkins formula has good diagnostic compatibility with the direct LDL-C measurement <sup>49</sup>
<b>Uganda</b>	HIV > 50 yrs of age	Direct LDL-C on C311 chemistry analyzer: <i>Roche Diagnostics</i>	Friedewald	2021	Friedewald equation reliably calculated LDL-C <sup>50</sup>
<b>United Kingdom</b>	Type II Diabetics; Statins	LDL-C immunoassay on Synchron DxC analyser: <i>Beckman-Coulter</i>	Friedewald, de Cordova	2015	De Cordova agreed more closely with directly measured LDL-C than Friedewald <sup>51</sup>

## References

1. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972;18(6):499-502.
2. DeLong DM, DeLong ER, Wood PD, et al. A comparison of methods for the estimation of plasma low- and very low-density lipoprotein cholesterol. The Lipid Research Clinics Prevalence Study. *JAMA* 1986;256(17):2372-7.
3. Rao A, Parker AH, el-Sheroni NA, et al. Calculation of low-density lipoprotein cholesterol with use of triglyceride/cholesterol ratios in lipoproteins compared with other calculation methods. *Clin Chem* 1988;34(12):2532-4.
4. Hattori Y, Suzuki M, Tsushima M, et al. Development of approximate formula for LDL-cholesterol, LDL-apo B and LDL-cholesterol/LDL-apo B as indices of hyperapobetalipoproteinemia and small dense LDL. *Atherosclerosis* 1998;138(2):289-99. doi: 10.1016/s0021-9150(98)00034-3
5. Puavilai W, Laoragpongse D. Is calculated LDL-C by using the new modified Friedewald equation better than the standard Friedewald equation? *J Med Assoc Thai* 2004;87(6):589-93.
6. Anandaraja S, Narang R, Godeswar R, et al. Low-density lipoprotein cholesterol estimation by a new formula in Indian population. *Int J Cardiol* 2005;102(1):117-20. doi: 10.1016/j.ijcard.2004.05.009
7. Teerakanchana T, Puavilai W, Suriyaprom K, et al. Comparative study of LDL-cholesterol levels in Thai patients by the direct method and using the Friedewald formula. *Southeast Asian J Trop Med Public Health* 2007;38(3):519-27.
8. Ahmadi SA, Boroumand MA, Gohari-Moghaddam K, et al. The impact of low serum triglyceride on LDL-cholesterol estimation. *Arch Iran Med* 2008;11(3):318-21. doi: 08113/AIM.0014
9. Chen Y, Zhang X, Pan B, et al. A modified formula for calculating low-density lipoprotein cholesterol values. *Lipids Health Dis* 2010;9:52. doi: 10.1186/1476-511X-9-52 [published Online First: 20100521]
10. Vujovic A, Kotur-Stevuljevic J, Spasic S, et al. Evaluation of different formulas for LDL-C calculation. *Lipids Health Dis* 2010;9:27. doi: 10.1186/1476-511X-9-27 [published Online First: 20100310]
11. de Cordova CM, de Cordova MM. A new accurate, simple formula for LDL-cholesterol estimation based on directly measured blood lipids from a large cohort. *Ann Clin Biochem* 2013;50(Pt 1):13-9. doi: 10.1258/acb.2012.011259 [published Online First: 2012/10/31]
12. Martin SS, Blaha MJ, Elshazly MB, et al. Comparison of a novel method vs the Friedewald equation for estimating low-density lipoprotein cholesterol levels from the standard lipid profile. *JAMA* 2013;310(19):2061-8. doi: 10.1001/jama.2013.280532
13. Sampson M, Ling C, Sun Q, et al. A New Equation for Calculation of Low-Density Lipoprotein Cholesterol in Patients With Normolipidemia and/or Hypertriglyceridemia. *JAMA Cardiol* 2020;5(5):540-48. doi: 10.1001/jamacardio.2020.0013
14. Dudi P, Vishwajeet, Yadav SRM, et al. Derivation of a Modified Friedewald's equation for LDL cholesterol estimation in Sub-Himalayan population: A comparative study. *J*

- Family Med Prim Care* 2022;11(5):1826-33. doi: 10.4103/jfmpc.jfmpc\_1063\_21 [published Online First: 20220514]
15. Sampson M, Wolska A, Meeusen JW, et al. Identification of Dysbetalipoproteinemia by an Enhanced Sampson-NIH Equation for Very Low-Density Lipoprotein-Cholesterol. *Front Genet* 2022;13:935257. doi: 10.3389/fgene.2022.935257 [published Online First: 20220714]
  16. Naser A, Isgandarov K, Güvenç TS, et al. Comparison of Novel Martin/Hopkins and Sampson Equations for Calculation of Low-Density Lipoprotein Cholesterol in Diabetic Patients. *Arq Bras Cardiol* 2022 doi: 10.36660/abc.20210641 [published Online First: 20220627]
  17. Shi B, Wang HY, Liu J, et al. Directly Measured vs. Calculated Low-Density Lipoprotein Cholesterol Does Not Identify Additional Individuals With Coronary Artery Disease and Diabetes at Higher Risk of Adverse Events: Insight From a Large Percutaneous Coronary Intervention Cohort in Asia. *Front Cardiovasc Med* 2022;9:932878. doi: 10.3389/fcvm.2022.932878 [published Online First: 20220707]
  18. Bauer F, Seibert FS, Rohn B, et al. Estimation of LDL cholesterol in chronic kidney disease. *Eur J Prev Cardiol* 2021;28(12):1402-08. doi: 10.1093/eurjpc/zwaa003
  19. Ephraim RKD, Acheampong E, Swaray SM, et al. Developing a Modified Low-Density Lipoprotein (M-LDL-C) Friedewald's Equation as a Substitute for Direct LDL-C Measure in a Ghanaian Population: A Comparative Study. *J Lipids* 2018;2018:7078409. doi: 10.1155/2018/7078409 [published Online First: 20181231]
  20. Reiber I, Mark L, Paragh G, et al. Comparison of low-density lipoprotein cholesterol level calculated using the modified Martin/Hopkins estimation or the Friedewald formula with direct homogeneous assay measured low-density lipoprotein cholesterol. *Arch Med Sci* 2022;18(3):577-86. doi: 10.5114/aoms.2020.97847 [published Online First: 20200803]
  21. Ramasamy J, Job V, Mani T, et al. Calculated values of serum LDL-cholesterol (LDL-C) - for better or worse? *Nutr Metab Cardiovasc Dis* 2021;31(5):1486-93. doi: 10.1016/j.numecd.2021.01.016 [published Online First: 20210129]
  22. Malve H, Asalkar A. Which LDL Value Should Clinicians Look at? *J Lab Physicians* 2021;13(2):129-33. doi: 10.1055/s-0041-1729131 [published Online First: 20210526]
  23. Tomo S, Sankanagoudar S, Shukla R, et al. Validation of a novel method for determination of low-density lipoprotein cholesterol levels in Indian patients with type 2 diabetes. *Diabetes Metab Syndr* 2022;16(4):102448. doi: 10.1016/j.dsx.2022.102448 [published Online First: 20220313]
  24. Atabi F, Mohammadi R. Clinical Validation of Eleven Formulas for Calculating LDL-C in Iran. *Iran J Pathol* 2020;15(4):261-67. doi: 10.30699/ijp.2020.110379.2174 [published Online First: 20200718]
  25. Molavi F, Namazi N, Asadi M, et al. Comparison common equations for LDL-C calculation with direct assay and developing a novel formula in Iranian children and adolescents: the CASPIAN V study. *Lipids Health Dis* 2020;19(1):129. doi: 10.1186/s12944-020-01306-7 [published Online First: 20200606]
  26. Egbaria A, Saliba W, Zafrir B. Assessment of Low LDL Cholesterol in Patients Treated by PCSK9 Inhibition: Comparison of Martin/Hopkins and Friedewald Estimations. *Cardiovasc Drugs Ther* 2021;35(4):787-92. doi: 10.1007/s10557-020-07107-5 [published Online First: 20201123]

27. Zafzir B, Saliba W, Flugelman MY. Comparison of Novel Equations for Estimating Low-Density Lipoprotein Cholesterol in Patients Undergoing Coronary Angiography. *J Atheroscler Thromb* 2020;27(12):1359-73. doi: 10.5551/jat.57133 [published Online First: 20200702]
28. Cicero AFG, Fogacci F, Patrono D, et al. Application of the Sampson equation to estimate LDL-C in children: Comparison with LDL direct measurement and Friedewald equation in the BLIP study. *Nutr Metab Cardiovasc Dis* 2021;31(6):1911-15. doi: 10.1016/j.numecd.2021.02.034 [published Online First: 20210319]
29. Sarzani R, Giuliotti F, Allevi M, et al. The calculation of LDL-cholesterol: Comparison of traditional and recent equations and their impact on lipid control rates in a hypertensive population at variable cardiovascular risk. *Atherosclerosis* 2021;331:e166.
30. Ouchi G, Komiya I, Taira S, et al. Triglyceride/low-density-lipoprotein cholesterol ratio is the most valuable predictor for increased small, dense LDL in type 2 diabetes patients. *Lipids Health Dis* 2022;21(1):4. doi: 10.1186/s12944-021-01612-8 [published Online First: 20220107]
31. Lee J, Jang S, Jeong H, et al. Validation of the Friedewald formula for estimating low density lipoprotein cholesterol: the Korea National Health and Nutrition Examination Survey, 2009 to 2011. *Korean J Intern Med* 2020;35(1):150-59. doi: 10.3904/kjim.2017.233 [published Online First: 20180321]
32. Song Y, Lee HS, Baik SJ, et al. Comparison of the effectiveness of Martin's equation, Friedewald's equation, and a Novel equation in low-density lipoprotein cholesterol estimation. *Sci Rep* 2021;11(1):13545. doi: 10.1038/s41598-021-92625-x [published Online First: 20210629]
33. Choi R, Park MJ, Oh Y, et al. Validation of multiple equations for estimating low-density lipoprotein cholesterol levels in Korean adults. *Lipids Health Dis* 2021;20(1):111. doi: 10.1186/s12944-021-01525-6 [published Online First: 20210920]
34. Chung S. Usefulness of the Martin Method for Estimation of Low-Density Lipoprotein Cholesterol in Coronary Atherosclerosis. *Med Princ Pract* 2018;27(1):8-14. doi: 10.1159/000485977 [published Online First: 20171204]
35. Barakett-Hamade V, Ghayad JP, McHantaf G, et al. Is Machine Learning-derived Low-Density Lipoprotein Cholesterol estimation more reliable than standard closed form equations? Insights from a laboratory database by comparison with a direct homogeneous assay. *Clin Chim Acta* 2021;519:220-26. doi: 10.1016/j.cca.2021.05.008 [published Online First: 20210513]
36. Orimadegun BE, Ogah F, Oyedele OB, et al. Plasma Low-Density Lipoprotein Cholesterol Estimated by Friedewald Compared to Martin-Hopkins Equation in Nigerian Population. *West Afr J Med* 2021;38(3):255-61.
37. Penson P, Martin S, Henney N, et al. Comparison of LDL-C calculation by friedewald and martin/hopkins methods in 12,243 adults from the United States of America. *European Heart Journal* 2020;41(Supplement\_2):ehaa946. 2932.
38. Ginsberg HN, Rosenson RS, Hovingh GK, et al. LDL-C calculated by Friedewald, Martin-Hopkins, or NIH equation 2 versus beta-quantification: pooled alirocumab trials. *J Lipid Res* 2022;63(1):100148. doi: 10.1016/j.jlr.2021.100148 [published Online First: 20211111]
39. Sajja A, Park J, Sathiyakumar V, et al. Comparison of Methods to Estimate Low-Density Lipoprotein Cholesterol in Patients With High Triglyceride Levels. *JAMA Netw Open*

- 2021;4(10):e2128817. doi: 10.1001/jamanetworkopen.2021.28817 [published Online First: 20211001]
40. Cao J, Remaley AT, Guan W, et al. Performance of novel low-density lipoprotein-cholesterol calculation methods in predicting clinical and subclinical atherosclerotic cardiovascular disease risk: The Multi-Ethnic Study of Atherosclerosis. *Atherosclerosis* 2021;327:1-4. doi: 10.1016/j.atherosclerosis.2021.04.018 [published Online First: 20210508]
41. Cartier LJ, St-Coeur S, Robin A, et al. Impact of the Martin/Hopkins modified equation for estimating LDL-C on lipid target attainment in a high risk patient population. *Clin Biochem* 2020;76:35-37. doi: 10.1016/j.clinbiochem.2019.12.002 [published Online First: 20191213]
42. Azimi V, Farnsworth CW, Roper SM. Comparison of the Friedewald equation with Martin and Sampson equations for estimating LDL cholesterol in hypertriglyceridemic adults. *Clin Biochem* 2022 doi: 10.1016/j.clinbiochem.2022.07.005 [published Online First: 20220726]
43. Rana MN, Kim CH, Sullivan CE, et al. Performance of Methods to Estimate Low-Density Lipoprotein Cholesterol in Women With and Without HIV Infection. *J Acquir Immune Defic Syndr* 2021;87(1):750-54. doi: 10.1097/QAI.0000000000002631
44. Schneider EE, Sarkar S, Margolick JB, et al. Short Communication: Comparison of Calculated Low-Density Lipoprotein Cholesterol (LDL-C) Values in HIV-Infected and HIV-Uninfected Men Using the Traditional Friedewald and the Novel Martin-Hopkins LDL-C Equations. *AIDS Res Hum Retroviruses* 2020;36(3):176-79. doi: 10.1089/AID.2019.0220 [published Online First: 20200106]
45. Gasko R. Comparison of LDL-C calculation by Martin, Sampson and old Friedewald methods in real data and synthetic data set. *Vnitr Lek* 2021;67(E-2):9-17.
46. Rossouw HM, Nagel SE, Pillay TS. Comparability of 11 different equations for estimating LDL cholesterol on different analysers. *Clin Chem Lab Med* 2021;59(12):1930-43. doi: 10.1515/cclm-2021-0747 [published Online First: 20210812]
47. Rerksuppaphol L, Rerksuppaphol S. Comparison of equations for the calculation of low-density lipoprotein cholesterol in thai population. *J Nat Sci Biol Med* 2021;12
48. Dinc Asarcikli L, Kis M, Guvenc TS, et al. Usefulness of novel Martin/Hopkins and Sampson equations over Friedewald equation in cardiology outpatients: A CVSCORE-TR substudy. *Int J Clin Pract* 2021;75(6):e14090. doi: 10.1111/ijcp.14090 [published Online First: 20210220]
49. Alpdemir M, Alpdemir MF. Comparison of Martin and Friedewald equation for estimated LDL-C in adults. *Turk Kardiyol Dern Ars* 2021;49(8):619-26. doi: 10.5543/tkda.2021.90446
50. Jude E, Brian S, Stephen K, et al. Expanding The Use Of Friedewald Equation In Lipid Profile Testing Among HIV Positive And Negative Elderly Persons. *medRxiv* 2021
51. Sathyapalan T, Atkin SL, Kilpatrick ES. Low density lipoprotein-cholesterol variability in patients with type 2 diabetes taking atorvastatin compared to simvastatin: justification for direct measurement? *Diabetes Obes Metab* 2010;12(6):540-4. doi: 10.1111/j.1463-1326.2009.01190.x