THE VALUE OF A SERUM AND ALBUMIN MIXTURE FOR USE IN THE DETECTION OF BLOOD GROUP ANTIGEN-ANTIBODY REACTIONS

BY

F. STRATTON AND E. R. DIMOND

From the Regional Blood Transfusion Service, Manchester

(RECEIVED FOR PUBLICATION JANUARY 25, 1955)

It has been suggested by Wiener and Hurst (1947) and Wiener, Hurst, and Sonn-Gordon (1947) that mixtures of bovine albumin and human plasma are more effective as diluents for the detection of incomplete antibodies than the widely employed 20% bovine albumin introduced by Diamond and Abelson (1945) or the human plasma alone used by Wiener (1945). This has been confirmed by other workers, and Mollison (1951) has recommended mixing a suspension of red cells in 20% albumin with the test serum diluted in group AB human plasma.

The present investigation was undertaken to determine whether the 20% bovine albumin test, which is open to criticism on a number of grounds, could be improved by substituting a mixture of human serum and bovine albumin for the diluent. Human plasma was not used because of clotting difficulties. After initial experiments it was decided to compare 20% bovine albumin with an equal parts mixture of human serum and 30% bovine albumin. Saline, and often human serum alone,

was also included in the comparisons which were made with respect to different blood group antigen-antibody reactions.

Materials and Methods

Red Cells.—Red cells were obtained in the form of clotted blood. The cells recovered from the clot were washed three times in normal saline and packed. The supernatant saline was completely removed from the packed cells. They were then resuspended at 10% strength in the appropriate diluting fluid. The test cells were of group O unless otherwise stated.

Human Serum.—The serum used for diluting the bovine albumin was obtained from donors of group AB and was either fresh or had been kept frozen solid for varying periods of time, and is referred to as AB serum. It was free from all blood group antibodies and had no rouleau-forming properties.

Antisera.—These human sera contained Rh or other blood group antibodies of various strengths and had been preserved frozen solid.

Bovine Albumin.—Armour's 30% bovine albumin (or bovine plasma) was used in the tests. The 20% albumin was made by diluting this with half its

TABLE I
RESULTS OF VARIOUS TESTS PERFORMED ON SERA CONTAINING WEAK ANTIBODIES (ANTI-D)

Serum No.	Tests																	
	Coombs Test			Papainized Red Cells		Agglutinins 37° C. in Saline			Titration of Serum in 20% Albumin × R ₁ r Cells at 37° C.					Titration of Serum in Serum-Albumin Mixture × R ₁ r Cells at 37° C.				
	R ₁ r	R ₂ r	rr	(R ₁ r + R ₂ r)*	rr	R ₁ r	R ₂ r	rr	1: 1	1: 2	1: 4	1: 8	Control	1: 1	1: 2	1: 4	1: 8	Control
16 A 14	++	++	—	++	—	—	—	—	—	—	—	—	—	+	—	—	—	—
16 A 43	+	+	—	++	—	—	—	—	—	—	—	—	—	+	w	—	—	—
20 F 15	++	+++	—	+++	—	—	w	—	w	—	—	—	—	++	+	w	—	—
20 F 12	—	+	—	+++	—	—	—	—	w	—	—	—	—	—	—	—	—	—
20 G 15	+++	+++	—	+++	—	—	—	—	—	—	—	—	—	—	—	—	—	—
20 G 44	++++	++++	—	+++	—	—	—	—	—	—	—	—	—	—	—	—	—	—

* Cell mixture.

Papainized Red Cells.—These were prepared and used as described by Stratton (1953).

Results were scored as follows:

++++ = Very strong agglutination with a few free cells.

+++ = Many large clumps of cells clearly visible with more free cells.

++ = Multiple clumps (10-12 cells) just visible to the naked eye, with many free cells.

+ = Small clumps (6-8 cells) visible microscopically; many free cells.

w=Clumps (3-4 cells) few in number;
many free cells.

— = Negative; completely free cell suspension.

Results

Detection of Weak Rh Antibodies.—Table I shows the results of testing sera containing weak anti-D antibodies by various methods. It will be observed that two sera negative in the 20% albumin test were weakly positive in the serum-albumin diluent, while two others giving very weak reactions in the former diluent gave stronger ones in the latter. All the antibodies in this table were weak but were detected using papainized cells and by the Coombs test. The last two sera in the table contained antibodies not demonstrated in the saline, 20% albumin or serum-albumin agglutination tests.

Enhancement of Titre of Rh-incomplete Antibodies by the Use of Serum-Albumin Mixture.—This enhancement of titre has been described before and was found to be usually two- to eight-fold. Fourteen incomplete anti-D sera of various titres were tested and all showed the greater effectiveness of serum-albumin in this respect. The control tests using Rh-negative cells all gave negative results. Table II illustrates a typical case.

Diluent				Dilutions							Control
				1:1	1:2	1:4	1:8	1:16	1:32	1:64	
20% albumin	..	..	..	++++	+++	++	+	—	—	—	—
Serum-albumin	..	..	..	++++	++++	+++	++	+	+	—	—
Saline	..	..	..	—	—	—	—	—	—	—	—
AB serum	..	..	..	+++	++	+	—	—	—	—	—

* Control titration against Rh-negative cells was negative throughout.

Occurrence of Zoning in Detection of Rh Antibodies.—Table III shows the titration of three anti-Rh sera with various Rh-positive cells in different diluents. All these sera showed zoning when the titration was performed using 20% albumin, but the zoning in each case was of a different kind and degree. In Table III (a), the zone using R_2r test cells was a marked one in the middle of the titration; it did not occur so markedly using type R_1R_1 (CDe/CDe) cells since the serum was of anti-C+D variety and zoning most commonly occurs only with respect to D. The serum in Table III (b) showed zoning starting earlier in the titration in 20% albumin. The third serum (Table III (c)) shows only a slight zone which is most marked in the lower dilutions, 1:1, 1:2, of the serum. It will be observed that the dilutions which gave negative results within the zones of the 20% albumin titrations gave unmistakable positive agglutination with serum-albumin. In the case of the second and third sera, if a single tube albumin test were used, the weak result at 1:1 might readily have passed unobserved, whereas with serum-albumin the corresponding result was + + +. Serum 20 F 30 (Table III (c)), being anti-D unlike the other two sera, showed a zone in 20% albumin against both R_1 and R_2 type cells.

Detection of Reaction between Incomplete Rh Antibodies and D^u Cells.—It is known from previous work (Stratton and Renton, 1949) that D^u cells of low grade are not agglutinated by incomplete Rh antibodies in a 20% albumin diluent. Table IV (a) shows the titration of an incomplete anti-D serum using various diluents and cells of type $R_1^u r$ (CD^ue/cde) low grade. Here the expected negative result was obtained using 20% albumin, but with serum-albumin the result was positive. It is interesting to note that this titration showed a zone. In Table IV (b) titration of the same serum with ordinary R_1r cells (CDe/cde) using the same diluents is shown. This serum was specially selected and known to be valuable in detecting D^u positive cells when using the indirect Coombs technique. When a number of other sera were used, some of lower titre, no agglutination was obtained even using serum-albumin diluent and CD^u/cde cells of low grade. Table IV (c) shows the results of titrating the same selected serum with cells of type $R_2^u r$ (cD^uE/cde) high grade. Here an agglutination was obtained in both 20% albumin and serum-albumin diluents, but the titre with the latter is much higher and the individual results with each dilution stronger. When a different, incomplete anti-D serum of

lower titre was titrated with these high grade D^u cells, a negative result was obtained using 20% bovine albumin, but a positive one using serum-albumin diluent (Table IV (d)).

Detection of Other Varieties of Rh Antibodies.—Other varieties of Rh antibodies were found to behave in a manner similar to the anti-D and anti-C+D antibodies which have been considered so far. An anti-c antibody (mainly incomplete) had a titre of 1 in 64 in 20% albumin, 1 in 256 in serum-albumin, and 1 in 4 in saline against Rh-negative (cde/cde) cells, and was completely negative with R_1R_1 (CDe/CDe) cells in these media. An anti-E antibody of titre 1 in 4 in albumin was of titre 1 in 16 in serum-albumin against R_2R_2 (cDE/cDE) cells and control titrations were again satisfactorily negative using R_1R_1 and Rh negative cells.

Detection of Complete Antibodies.—Two varieties of complete antibodies were examined; first, what was considered to be a serum containing largely complete Rh antibodies, in view of the fact that its titre was the same in saline, 20% albumin, and whole serum, namely, 1 in 32. The titre in serum-albumin mixture was 64.

Anti-A and anti-B antibodies, some of which had been selected as being of low titre in saline, were titrated out in various diluents against cells of appropriate group. There was no evident enhancement of agglutination using either 20% albumin or serum-albumin mixture and, indeed, with some sera the results were worse than when a saline diluent was used.

Anti-Kell.—Six anti-Kell sera were titrated with known Kell-positive cells in various diluents; the results of a typical titration are shown in Table V. It is interesting to observe the negative

TABLE V
TITRATION OF AN ANTI-KELL SERUM (NO. 13 F 1) WITH KELL-POSITIVE CELLS AT 37° C. IN VARIOUS DILUENTS

Diluent	Dilutions					Control
	1:1	1:2	1:4	1:8	1:16	
20% albumin ..	—	—	—	—	—	—
Serum-albumin ..	++	+	w	—	—	—
Saline ..	—	—	—	—	—	—
AB serum ..	+++	++	++	+	—	—

reaction in the 20% albumin and the positive result which was obtained in the serum-albumin titration; also an even stronger agglutination occurred in serum alone. Grove-Rasmussen (personal communication) informed us he had found this using several different anti-Kell sera.

Five out of six anti-Kell sera we examined showed this higher titre in whole serum than in any other diluent.

Anti-Duffy, Anti-S, and Anti-Kidd.—Several examples of each of these sera were tested with the various diluents, but all gave negative results with cells containing the agglutinin for these antisera.

Anti-Lewis.—Various tests have been made titrating anti-Le^a and anti-Le^b antibodies with the appropriate cells at 16° C. and 37° C., using the usual diluents. The results showed little enhancement of agglutination when 20% albumin or serum-albumin was used, compared with saline, but it is probably advantageous to use an albumin technique at 37° C. if it is suspected that incomplete Lewis antibodies are present in a serum.

Effect of Albumin Mixture on Activity of Cold Agglutinins.—Table VI shows the titration of a non-specific cold antibody at 8° C. and 37° C. There is little difference between the titres and strengths of agglutination using 20% albumin and serum-albumin, but both give a higher titre than saline.

TABLE VI

TITRATION OF SERUM (NO. 12 D 17) CONTAINING A NON-SPECIFIC COLD AGGLUTININ WITH Rh-NEGATIVE CELLS AT 8° C. AND 37° C. IN VARIOUS DILUENTS

Diluent	Dilutions						Control
	1:1	1:2	1:4	1:8	1:16	1:32	
(a) 8° C.: 20% albumin	++++	+++	+++	++	w	—	—
Serum-albumin	++++	+++	+++	++	+	—	—
Saline ..	+++	++	+	+	—	—	—
(b) 37° C.: 20% albumin	—	—	—	—	—	—	—
Serum-albumin	—	—	—	—	—	—	—
Saline ..	—	—	—	—	—	—	—

Centrifugation Experiments.—Experiments were carried out to see whether time could be saved during the performance of albumin tests by centrifuging the tubes following a limited incubation period and then reading the tests. An anti-D serum titrated in both 20% albumin and serum-albumin mixture was employed. The tubes were incubated for 20 minutes and then centrifuged at low speed (500 r.p.m.) for five minutes and the results read. Comparison with the standard method showed that no serious error would result from using this method if a result is required quickly. Centrifugation experiments have been conducted with only a limited number of anti-Rh sera and not with the full series of tests.

Commentary

The albumin test, which is widely used for routine purposes, suffers from certain defects. Some of these can be remedied by using serum-albumin mixture in the proportions suggested instead of the 20% albumin mixture normally used.

It has been shown (Stratton, 1953) that as many as 5 to 7% of weak Rh antibodies in the maternal serum might be missed if the 20% bovine albumin test were used alone; when a serum-albumin mixture is used many fewer Rh antibodies would remain undetected.

All serological techniques for the detection of blood group antigen-antibody reactions are liable to give false negative results due to the occurrence of zoning, but the albumin test is particularly prone to this phenomenon. This means that when using a 20% bovine albumin diluent a single tube test cannot safely be employed and the test should be performed in the form of a titration using up to six tubes. The results reported here show that a serum-albumin diluent reduces the danger of these false negative results. Even where the first tube of a serum titration shows a weak positive result, due to zoning when 20% bovine-albumin is used, the substitution of the serum-albumin mixture would result in a stronger positive result. It should not be assumed, however, that if a serum-albumin mixture is used every difficulty due to zoning will be eliminated in this test, but false negative results due to this cause will be considerably diminished.

Stratton and Renton (1949) showed that the D^u type of cells, especially if of low grade, will not react with incomplete anti-D sera in 20% bovine albumin, but if the D^u type of cell is of high grade occasional positive reactions will be obtained depending upon the incomplete anti-D serum that is used. When a serum-albumin diluent is used depending upon the particular anti-D sera, some low grade D^u positive cells may be detected; even so, a zone may occur in these titrations. Nevertheless, it would seem that, in a population in which there occurs a large percentage of the D^u-positive type of person and where the albumin test is used as a screening test in the first instance for the detection of D-positive persons, materially better results would be obtained if a serum-albumin mixture were used for the original screening test. It should be emphasized that negative results would need to be retested using the Coombs test. One should not attempt to detect low-grade D^u cells using the albumin technique alone, whatever diluent is used.

Rarely, Rh antibodies are not detectable by the antiglobulin or other tests, but only using the albumin technique. If these Rh antibodies are detected in the serum of an Rh-negative mother during the antenatal period, the subsequent sample usually gives a positive indirect Coombs test. Anti-E antibody is one that is most often preferentially detected by the use of the albumin technique.

Kell is an antigen to which persons readily become immunized and anti-Kell antibodies are often detected in human sera. The Coombs test has most often been used for the detection of these antibodies, and the 20% bovine albumin test does not usually give a positive result. Our results suggest, however, that if serum-albumin is used positive results will be obtained. Grove-Rasmussen (personal communication) suggested that good results were obtained if a serum diluent were used, and this has been confirmed.

Anti-Duffy, anti-Kidd, and sometimes anti-S cannot be detected at all using the albumin technique, whichever diluent is used, and the chances of detecting Lewis antibodies are not materially increased by the substitution of a serum-albumin mixture for a 20% one.

In the case of anti-A and anti-B saline agglutinins, our impression, after testing many samples, was that these agglutinins are not as readily detected in albumin or in serum-albumin mixtures as they are in saline, although they would be unlikely to give completely negative results in the first two diluents and positive ones in saline.

Experiments were carried out to determine whether the increased rate of sedimentation of the red cells was the sole reason for the enhanced agglutination of the serum-albumin mixture as compared with 20% albumin. These experiments consisted of reading results of parallel titrations after incubation periods ranging from one to four hours. There was no evidence to support this, since in all cases the serum-albumin results were stronger than those where 20% albumin was used. These experiments did suggest that economies in time, such as reading the test after one and a half hours' incubation, would not have such serious effects in preventing the detection of positive results if serum-albumin were used as a diluent compared with 20% bovine albumin. It seems to us, therefore, that the time of incubation might be reduced accordingly when serum-albumin mixture is used. Low-speed centrifugation to reduce the time of incubation was also considered and an

incubation time of 20 minutes at 37° C. in the water-bath followed by low-speed centrifugation (500 r.p.m.) might well be used where speed is desired. Our tests with this technique have only been concerned with a limited number of Rh antibodies.

Many observers have shown that other substances will act in the same way as bovine albumin and human plasma or serum. Gelatin and dextran, for example, have been employed, and one of us found, of many substances tried, that methyl cellulose was the best. All these substances suffered from the defect that they tend to cause false positive results which make the interpretation of the test most difficult. The selection of a suitable test medium, therefore, involves finding one which will not give false positive results but will detect blood group antigen-antibody reactions with the minimum of difficulty.

In the authors' experience it is an advantage to add human serum in the proportions suggested to bovine albumin, and such a 50/50 mixture will not increase the false positive results or the difficulty of reading the test. The serum used in making the mixture should be free from blood-group antibodies and all rouleau-forming properties, and we are in favour of freezing it solid at least a week or ten days before making up the mixture. After mixing it is stored at 4° C. and not kept for more than a few days.

The reading and interpretation of the albumin test have always presented difficulty. The cell deposit should be carefully removed from the tube and spread upon a microscope slide. At this stage aggregation of the red cells is liable to be present, especially after centrifugation, and gives the effect of comets, i.e., a clump of cells from which a tail of cells can be seen trailing away. It is important to leave the slide a few minutes on the bench before placing it under the low power of the microscope and then to examine a moving film. This is done by tilting the slide on the stage when the break-up of the cell aggregates will be observed, the true agglutinates remaining intact. In the case of saline agglutination tests, it is often possible to distinguish rouleau from true agglutination microscopically, but in the case of albumin or serum-albumin tests it is quite impossible to make this distinction.

The albumin technique is widely employed in the crossmatching test and often in the form of a single tube test. While the authors would not

recommend this as a routine method of cross-matching, nevertheless it does seem a considerable advantage to use the serum-albumin mixture suggested in place of 20% bovine albumin in such a test, and such a mixture would also be advantageous if the test were done in the form of a titration. In the crossmatching test, however, it seems preferable to use a saline agglutination method in tubes at room temperature and at 37° C. coupled with the use of the antiglobulin technique.

REFERENCES

- Diamond, L. K., and Abelson, N. M. (1945). *J. Lab. clin. Med.*, **30**, 204 and 668.
- Grove-Rasmussen, M. Personal communication.
- Mollison, P. L. (1951). *Blood Transfusion in Clinical Medicine*. Blackwell, Oxford.
- Mourant, A. E., and Race, R. R. (1952). *The Rh Blood Groups and Their Clinical Effects*. Medical Research Council Memorandum No. 27. H.M.S.O., London.
- Stratton, F. (1953). *Lancet*, **1**, 1169.
- and Renton, P. H. (1949). *Brit. med. J.*, **2**, 682.
- Wiener, A. S. (1945). *J. Lab. clin. Med.*, **30**, 662.
- and Hurst, J. G. (1947). *Exp. Med. Surg.*, **5**, 285.
- — and Sonn-Gordon, E. B. (1947). *J. exp. Med.*, **86**, 267.