Renal necrosis

J. F. SMITH

From the Department of Morbid Anatomy, University College Hospital Medical School, London

Renal necrosis due to ischaemia with a brief mention of lesions produced by some chemical poisons, particularly those which may act on renal blood vessels and thereby have an ischaemic action, will be the main considerations in this article. Such causes of renal necrosis have no clinical relevance even if they exist: there is no evidence, for example, of acute or chronic lesions in the kidney due to carbon monoxide or cyanide poisoning or nitrous oxide anaesthetic accidents comparable to those seen in the brain. It is probable that the kidney epithelial cells, even those of the metabolically active proximal tubule, can survive severe hypoxia although their function may be impaired.

In assessing the effects of ischaemia it is necessary to recollect a few anatomical facts: the branches of the renal arteries from the arcuates onwards are virtual end arteries and the infarct following occlusion of one is familiar to all from early studies in pathology. It is easy to forget, however, that such infarcts often spare the medullary pyramids whose blood supply is derived from the vasa recta, except for their papillae which are supplied also by a pericalyceal artery (Baker, 1959). The fact that scars of infarcts following arterial emboli spare the medulla and thereby do not cause calyceal deformity is a further illustration of this fact (Smith, 1962) and useful in differential diagnosis.

Ischaemic cortical necrosis can vary in degree from focal, through minor and patchy, to gross as documented in the thorough study by Sheehan and Moore (1952) of lesions associated with pregnancy. In all there is some degree of tubular necrosis, and in all except the minor, some degree of vascular occlusion. Whether the vascular occlusion develops from an initial intravascular coagulation or as the result of arterial and arteriolar spasm is undecided and need not concern us here. However, the association with tubular damage and in particular with proximal tubular damage should be stressed. The effects of such occlusion can be contrasted with that seen in the kidney lesion associated with excessive excretion of pigment as in incompatible transfusion, the crush syndrome and what was at the time referred to as the kidney of septic abortion, now known to

have been the consequence of intravascular haemolysis following the entrance of soap solution into the vessels from the placental site. We now know that the renal tubular necrosis in such conditions is not a lower nephron necrosis, the term invented by Balduin Lucké (1946) after the initial observations of Dunn *et al*, in 1941, but it is certainly not a predominantly proximal tubular necrosis as occurs in ischaemia.

Perhaps it is as well to look at this matter in an historical way. Until 1940 most pathologists had accepted that the oliguria following incompatible blood transfusion was the result of the blockage of tubules by pigment casts. This hypothesis was given experimental support by the work of Baker and Dodds (1925). When he examined the kidneys of those who had died from renal failure following crushing injuries of the limbs Shaw Dunn (Dunn *et al*, 1941) was impressed not only by the myohaemoglobin pigment casts in the lumina of tubules but by damage to tubular epithelium and focal disruption of epithelium. He thought that the damage was in the broad ascending limb of Henle and he was influenced in this view by an experimental uric acid lesion he had studied in rabbits with Polson (Dunn and Polson, 1926). In looking at those old experiments, which were the foundation stone of the concept of lower nephron nephrosis, one must remember two things. In the first place the dose of uric acid was high (2 g/kg) and it is probable that its action on the tubular epithelium was a mechanical traumatic effect as it came out of solution, rather than a cytotoxic lesion. Secondly, although Shaw Dunn took a lot of trouble to localize the experimental lesion to the ascending limb of Henle he was not correct in so doing, as the microdissection studies carried out by Smith and Lee (1957) show that the damage is actually in the collecting tubules.

There is no experimental support for the concept of a 'lower nephron nephrosis' and it is largely abandoned. It is, however, in my view unfortunate that the concept of 'shock kidney' has taken its place (Oliver *et al*, 1951) which has its bedrock in the demonstration by the technique of microdissection that the lesions are scattered in the cortex and

medulla and the view that such scattered lesions in the kidney and other organs are characteristic of ischaemia consequent to shock. However, there is no evidence that shock as such causes a patchy ischaemia and considerable evidence that ischaemia causes more severe damage to proximal cortical tubules than elsewhere. I subscribe to the Bywaters doctrine (1948) that in the great majority of cases of renal failure following excessive pigment secretion are the result of a raised interstitial pressure following tubular disruption at points of sheering stress in nephrons. The stress is produced by lumina being blocked by pigment casts and damaged epithelium in lower parts of scattered nephrons.

In cases where there has been selective necrosis of proximal tubular epithelium the possibility of an ischaemic aetiology is more likely. Although this may happen following cardiac arrest or a fall in blood pressure, it is rare. When observed one must always enquire carefully into the possibility of a tubular poison. It is the characteristic lesion of inorganic mercury poisoning, fortunately now rare, but may occur with some antibiotics, and in experimental animals can be produced by paracetamol.

An important practical point in the diagnosis of tubular necrosis, particularly in human disease, is that in histological sections one may find the regenerative stage. It is important to recognize that epithelium with flattened basophilic cells with large nuclei and sometimes with nuclear crowding is indicative of regeneration following necrosis. Necrotic cells may no longer be present and even the stage of mitotic activity may have passed.

In considering necrosis of the renal medulla it is noteworthy that the florid type complicating severe pyelonephritis in diabetics or following obstruction of the lower urinary tract now attracts less attention than the papillary necrosis associated with analgesic nephropathy. The exact pathogenesis of both forms is still uncertain but it is likely that interference with the blood supply is the important factor. In the florid lesion in diabetics it is postulated that the exudate in the intermediate zone compresses the vasa recta and with loss of the major blood supply to the medulla extensive necrosis follows. The naked-eye appearance is that of a typical anaemic infarct which involves much of the pyramid, is not shrunken, and therefore easily detected.

In the kidney of analgesic nephropathy the ischaemic process is probably slower, the necrotic lesions have often shrunk by the time of necropsy and may appear as shrivelled, greenish-brown structures. Kincaid-Smith (1967) has drawn attention to the association of atrophic cortex over such shrunken non-functional pyramids and to the fact that the resultant cortical scars may alternate with areas of bulging hypertrophy where the cortical tissue can still drain effectively into the superolateral and non-necrotic parts of the pyramids. The experimental work she has reported with other colleagues (Kincaid-Smith *et al*, 1968) indicates that the ischaemic necrosis here may be due to a toxic action of the analgesics on the walls of the vasa recta. However, it must be recognized that the snag in accepting both forms of medullary necrosis as ischaemic is the absence of recent or organized thrombus in the vasa recta in nearly all cases.

References

- Baker, S. B. de C. (1959). The blood supply of the renal papilla. *British Journal of Urology*, **31**, 53-59.
- Baker, S. L., and Dodds, E. C. (1925). Obstruction of the renal tubules during the excretion of haemoglobin. *British Journal of Experimental Pathology*, **6**, 247-260.
- Bywaters, E. G. L. (1948). "Renal anoxia" (Letter). *Lancet*, **301**.
- Dunn, J. S., Gillespie, M., and Niven, J. S. F. (1941). Renal lesions in two cases of crush syndrome. *Lancet*, **2**, 549-552.
- Dunn, J. S., and Polson, C. J. (1926). Experimental uric acid nephritis. *Journal of Pathology and Bacteriology*, **29**, 337-352.
- Kincaid-Smith, P. (1967). Pathogenesis of the renal lesion associated with the abuse of analgesics. *Lancet*, **1**, 859-862.
- Kincaid-Smith, P., Saker, B. M., and McKenzie, I. F. C. (1968). Lesions in the vasa recta in experimental analgesic nephropathy. *Lancet*, **1**, 24.
- Lucké, B. (1946). Lower nephron nephrosis. *Military Surgeon*, **99**, 371-396.
- Oliver, J., MacDowell, M., and Tracy, A. (1951). The pathogenesis of acute renal failure associated with traumatic and toxic injury: renal ischemia, nephrotoxic damage and the ischemic episode. *Journal of Clinical Investigation*, **30**, 1307-1439.
- Sheehan, H. L., and Moore, H. C. (1952). *Renal Cortical Necrosis and the Kidney of Concealed Accidental Haemorrhage*. Blackwell, Oxford.
- Smith, J. F. (1962). The diagnosis of the scars of chronic pyelonephritis. *Journal of Clinical Pathology*, **15**, 522-526.
- Smith, J. F., and Lee, Y. C. (1957). Experimental uric acid nephritis in the rabbit. *Journal of Experimental Medicine*, **105**, 615-622.