The cells

Fibroblasts

M. ABERCROMBIE

From the Strangeways Research Laboratory, Cambridge

The fibroblast, or more cautiously the fibroblast-like cell, is the standard cell of the tissue culturist. Most of our knowledge of fibroblasts concerns what they do in culture. It is with cultured fibroblasts that I shall mainly deal. Questions of the taxonomy of fibroblasts I will leave till later. For the present I will simply assume that there is a roughly delimited class of cells observed *in vitro* that approximately corresponds to a roughly delimited class of cells observed in connective tissue, both classes being called fibroblasts.

Shape

A cell is identified as a fibroblast in culture partly by its shape, observed if possible over a period of time, and partly by its spatial relations to other similar cells. On the usual flat tissue culture substratum an isolated fibroblast, fresh from the organism, has the well-known elongated flattened form, oblong or triangular. In the case of an embryonic bird heart fibroblast it may be 50 to 100 μm long, up to 30 μm at its broadest, and only about 3 μm at its thickest (Fig. 1). Its environment in culture will certainly induce it to move, so it has a front edge that is convex in shape as seen in plan, somewhat irregular in outline, and less than 0.5 μm thick. Other parts of the outline may be similarly convex and irregular, but much of it, especially the lateral edges, is thicker, smoother, and often slightly concave. So long as it remains isolated it will hold approximately to this form, merely altering its dimensions and occasionally changing the position of its front edge. A macrophage can at first glance imitate a fibroblast but its changes of form are more rapid and radical.

How can one explain the fibroblast's shape? Essentially, by the disposition of its adhesions to the solid substratum. A fibroblast has no detectable internal skeleton that by its rigidity maintains the characteristic shape. The moment its adhesions are released it starts to draw itself into an approximately spherical form. Though the adhesions are of course changing, since the cell is moving, they maintain

their general pattern and can be said to pin the cell out on the rigid surface of the culture vessel. If the cell is not on a flat surface but is, say, adhering to the fibrils of a collagen lattice it will have a different shape—not flattened but cylindrical, with a very narrow leading edge.

Fibroblasts, however, are usually not isolated. They make contact with their neighbours. These contacts are adhesions, usually rather limited in extent, and they distort the form of the isolated cell. They link fibroblasts into the characteristic meshwork, which is one of the ways they are recognised to be fibroblasts. Each adhesion usually produces a smoothly contoured, often concave, bordered projection of the cell to meet its neighbour. These smooth, slightly concave borders of so much of the fibroblast, as seen in plan, are expressions of the fact that the cell is under considerable tension between its adhesions to substratum or to neighbouring cell. The tensions are very probably developed by an actin-myosin system.

Fig. 1 Phase-contrast photograph of living chick heart fibroblast. The cell has been moving to the right and is in the act of changing direction to move upwards in the picture. Bar = 20 μm .

Contractility

The actin-myosin system of fibroblasts is most strikingly displayed in the bundles of microfilaments that have been demonstrated in recent years by the application of fluorescent antibody prepared against the various proteins found in the myofibril of striated muscle (Goldman *et al.*, 1975). In the fibroblast these bundles, up to 1 μm thick and several μm long, lie roughly parallel to the substratum and are commonly aligned in the direction the cell is moving (Fig. 2). They have a remarkably complex make-up, containing proteins closely similar to the actin, myosin, tropomyosin, α -actinin, and M-protein of striped muscle. No doubt more will be discovered. Not surprisingly, when the difficult feat of isolating these bundles of microfilaments was performed by Isenberg *et al.* (1976) they proved to be contractile when given ATP.

The bundles have some important differences from the myofibrils of striped muscle, which must be kept in mind. The individual proteins are not quite the same. The fibroblast, for example, has three kinds of actin (Rubenstein and Spudich, 1977), the main one being apparently absent from muscle tissue. The bundles contain extra proteins, such as filamin (Wang *et al.*, 1975), that are not found or are not conspicuous in myofibrils. And, most important of all, the bundles are transitory structures, appearing and sometimes disappearing in a matter of seconds. Furthermore, there is good reason to suppose that they are not the only components of the fibroblast contractile system, although they may be the most powerful. The general cytoplasmic matrix, especially at the leading end of a moving fibroblast, appears also to be contractile.

The contractile bundles of microfilaments have an important relation to the cell surface and to the substratum. One end of each bundle is attached to the cell surface (sometimes both ends) at a plaque of dense material just within the plasma membrane (Abercrombie *et al.*, 1971). Where these attachments to the membrane occur the membrane itself forms a localised and very tight adhesion to the substratum (Fig. 3) (Izzard and Lochner, 1976). A similar relation of bundles, plaques, and adhesions occurs at the cell-to-cell contacts (Heaysman and Pegrum, 1973). If now the substratum is not rigid like a culture dish but flexible like a collagen or fibrin substratum in the organism contraction of bundles throughout a meshwork of fibroblasts may produce deformations of whole areas of tissue. An example of this *in vivo* is the strange phenomenon of wound contraction, which in favourable circumstance may reduce the area of a skin wound to 10% of its original size. The evidence 20 years ago pointed

to contraction by the cells of the granulation tissue as the source of the motive force (Abercrombie *et al.*, 1956), and there is now further evidence for such activity by the population of fibroblasts (Gabbiani and Montandon, 1977). The role of tissue deformation by fibroblast contractility, especially in embryonic development, may well have unsuspected importance.

The fibroblast must therefore be regarded as a contractile cell. An appreciable part of its resources

Fig. 2 High voltage (800 kV) electron micrograph of 1.5- μm thick section of chick heart fibroblast. Several bundles of microfilaments are visible in the leading part of the cell (top right). Bar = 10 μm . (Courtesy of Mr J. P. Heath.)

Fig. 3 Interference reflection micrograph of chick heart fibroblast (same cell as Fig. 1 5 min later). Localised adhesions to the substratum appear as black streaks. Bar = 20 μm .

Fibroblasts

are applied to this function. Actin is the protein present in largest amount in the fibroblast in culture, forming up to 14% of the cell's proteins (Bray and Thomas, 1975). Fibroblasts *in vitro* are capable of exerting a force about one-tenth of that exerted by smooth muscle of the same cross-sectional area. Of course fibroblasts are not unique among non-muscle cells in their contractile proteins and contractile powers, but they seem to be among the leaders in this field.

Locomotion

The contractile proteins, and especially the bundles of microfilaments, no doubt play a part in fibroblast locomotion. In a migrating fibroblast the focal contacts from which the bundles spring are mostly near the front end of the cell. The bundles run backwards from their plaques, most of them disappearing into the microfilament system near the nucleus. The contraction of these bundles is well placed for dragging the main body of the cell forwards towards the anterior adhesions. Such contraction would produce only very limited locomotion—one step, so to speak. But the front edge, which differs from most of the periphery in appearance, does so because it is a region where the cell actively protrudes material forwards. This extension forms new adhesions to the substratum with new bundles of microfilaments attached. The old adhesions and their bundles fade out as the cell moves forwards past them. It is by such a continuous cycle of forming and losing adhesions that, on present evidence, we suppose locomotion takes place (Abercrombie *et al.*, 1977).

The machinery is capable of generating only very slow speeds, up to 100 μm per hour on a plane substratum, and an isolated cell will not follow a straight path for long, so it achieves a very small rate of displacement. Nevertheless, a *population* of fibroblasts can move with considerable efficiency in occupying space on a substratum that is devoid of other fibroblasts. This is because of a form of social behaviour termed contact inhibition. The region of extension of the cell located at its front end usually ceases to work when it touches another fibroblast and becomes instead a temporary adhesion. Hence the meshwork of adhering cells that is the usual form of a fibroblast population. Some other part of the cell periphery, free of contact, will now become a new front edge, as if released from competitive inhibition by the demise of the original front edge. A cell will therefore tend to move away from contacts, and, in aggregate, this can produce a directed streaming of fibroblasts into unoccupied territory.

Multiplication

It has always been very easy to get fibroblasts to multiply *in vitro*—in fact, they are commonly accused of swamping the growth of other kinds of cells in a culture. To multiply they must be provided with a rigid substratum of a size that enables them to spread to something like their usual length (anchorage-dependence, Stoker *et al.*, 1968). If that condition is met the media regularly used in tissue culture (in early days plasma and embryo extract, latterly serum) strongly encourage fibroblasts to go through the replicative cycle. It is becoming a little clearer where the stimulatory activity of serum resides. It seems to be in growth-promoting activity liberated by platelets when the serum is prepared. It may well be identified with the polypeptide, fibroblast growth factor, which Gospodarowicz isolated from pituitary (Gospodarowicz and Moran, 1975). This is not strictly specific for fibroblasts since it promotes growth also in smooth muscle, cartilage, and endothelium. Two other growth promoting polypeptides for which the fibroblast has receptors, EGF (Carpenter and Cohen, 1976) and MSA (Rechler *et al.*, 1977), have also been isolated, originally from outlandish sources.

Cultured fibroblasts generally drop out of the cell cycle, or cycle infrequently, when density of cell population becomes high. This can be observed in areas of high and low density within the same population in a culture, where the bulk medium is clearly able to support the more rapid cycling of the low density areas. This effect of density has always seemed a particularly interesting example of how the rapid cycling inducible by medium components can be shut off by a feed-back mechanism. There has been for long an oscillation between the theory that this density-dependent inhibition depends on the amount of cell-to-cell contact and the theory that it depends on local cell-induced changes in the concentration of medium components. Stoker (1973) has much advanced the claims of the second theory with evidence that intense stirring of the medium can produce an effect on the local multiplication rate.

The existence of the pronounced density-dependence of the cell cycle frequency *in vitro* has failed to stimulate investigations to discover whether the same holds *in vivo*. An exception is the report by Summerbell and Wolpert (1972) which showed that it did during the development of the chick limb.

A well known but still deeply puzzling outcome of the study of fibroblast growth *in vitro* is the Hayflick effect (Hayflick and Moorhead, 1961) whereby the number of replicative cycles that a fibroblast and its progeny can perform is limited to something like 60 unless there is transformation into

heteroploid and usually neoplastic form, a rare event in most species. The number of possible generations declines with the age of the donor; so whatever mechanism is at work operates *in vivo* as well as *in vitro*. The end of the period of multiplication *in vitro* is brought about by the decline and death of the cell population. A recent study by Holliday *et al.* (1977) has suggested that, in one respect, Hayflick's discovery is an artefact of culture. There seems to be *in vivo* a small class of stem cells capable of indefinite multiplication but continually giving rise to a proportion of progeny with a replicative life limited to 60 or so generations. The transplantation methods of tissue culturists have a very high probability of eliminating these rare stem cells and so limiting the life span of the cultured population. These are not, of course, stem cells of the usual kind producing a proportion of non-replicative progeny, as in many epithelial and in haematopoietic populations. Such progeny seem to be unknown amongst fibroblast populations.

Export

Later papers review the characteristic major exports of fibroblasts, collagen (Green *et al.*, 1966), the proteoglycans, and the enzymes that modify the extracellular substances. The production of these materials is, of course, very active in fibroblasts growing in culture (Manner, 1971; Steinberg, 1973), though the *proportion* of total protein synthesis devoted to them may be much less than *in vivo* (Schwarz *et al.*, 1976). The range of exports from a fibroblast, as from other cells, is probably greatly underestimated by present knowledge. An export that has until recently escaped detection is the LETS protein. This is a very large glycoprotein, identified by several groups and hence given several different names. It is exported on to the surface of fibroblasts in considerable amounts—up to 3% of the total fibroblast protein can be LETS (Yamada *et al.*, 1977). Other cell types produce smaller quantities. Most LETS remains closely associated with the cell surface and is indeed implicated in cell adhesion, but some moves further afield and appears in the culture medium or (as cold-insoluble globulin) in the blood plasma *in vivo*.

Some of the substances exported from a fibroblast are likely to be concerned with co-ordinating activities in the population in which it resides. For instance, fibroblasts produce substances that are growth-promoting for other fibroblasts (Millis *et al.*, 1977). In this connection the discovery, largely explored by Loewenstein, of communication channels through cell contacts is especially interesting. Direct export of small molecules (Pitts and Simms,

1977) occurs from cell to cell through gap junctions (Gilula *et al.*, 1972), though what role such transfers may have, other than making the population tend to metabolic uniformity, is not yet clear.

Activation

A general point must be made about the four activities of fibroblasts that I have chosen to discuss. Directed locomotion, active multiplication, the generation in a population of contractile force, and elaboration of extracellular material with relatively high tensile strength are much in evidence in culture and they are also the major features of the fibroblast's central function in repair processes. This is no accident. The fibroblasts that have been so much studied *in vitro* have almost always been activated by their preparation and subsequent conditions, in much the same way that fibroblasts in the organism are activated by injury. They are rarely allowed to settle down into the relatively quiescent state of normal connective tissue. Little of the information we have concerns the latter.

Taxonomy

My discussion so far has proceeded as if fibroblasts *in vitro* were a homogeneous and delimited type of cell with an *in-vivo* equivalent. I must now attempt to discuss the problems of identity entailed in writing about fibroblasts.

There is an *in-vitro* fibroblast phenotype. It is of considerable variability but, on the whole, is distinguishable from the other major types of cell. How are these fibroblasts related to the different kinds of cell in the tissue of origin? I am here considering the cells of primary or at all events short-term cultures that have not undergone prolonged selection and alteration *in vitro*. Probably the *in-vitro* fibroblasts are derived from cells that would be called fibroblasts *in vivo*, but there may be other cells, especially dissociated vascular endothelial cells, mixed in (see, for instance, Papayannopoulou and Martin, 1967). Furthermore, the *in-vitro* fibroblasts are most unlikely to be a random sample of the *in-vivo* cells. Setting up a culture, especially of adult cells, is often an occasion for great slaughter, as well as for differential growth of the survivors. Franks and Wilson (1977) have suggested that ultimately the only cells that survive are derived from the perivascular mesenchyme and from vascular endothelial cells.

Are the cells that emerge as tissue-culture fibroblasts equivalent, regardless of what organ they come from? Unquestionably they are not, and this is not just a matter of different proportions of the *in-vivo*

cell types contributing to the culture. *In-vitro* fibroblasts often look different when they come from different organs and are cultured under the same conditions. Fazzari (1930) studied the question with different explants placed side-by-side in the same culture. Though they become more alike with continued culture he thought that they could always be distinguished. Conrad and Hart (1973) thought the same. An embryologist would indeed be surprised if organ specificity of some kind was not detectable (Lewis and Wolpert, 1976). For instance, that the early mesenchyme has different properties in different parts of the embryo is shown by its influence on the differentiation of various epithelia (see Wessells, 1977). The fibroblasts of the tissue colturist is evidently a genus of many different species. Within any one of these species there may, of course, be much inherited variation between individual cells but this has not yet been much explored.

How firmly is the fibroblast committed to remaining a fibroblast and to reproducing only fibroblasts? Probably we can now discount early claims that in culture the fibroblast turned readily into a macrophage. It seems to be highly stable *in vitro*: it can coexist and multiply in the same culture with cartilage cells or myoblasts and remain distinct. But *in-vivo* at least some species of fibroblast undoubtedly are able under appropriate conditions to switch to other stable states, certainly to cartilage and bone. I do not think this necessarily means that there are uncommitted elements among what we call fibroblasts, or that fibroblasts are any less stable than cartilage cells or bone cells. But it does mean that the state of determination of the family of fibroblast, cartilage, bone, and perhaps other related cell types is an interesting question.

This work is supported by the Medical Research Council.

References

- Abercrombie, M., Dunn, G. A., and Heath, J. P. (1977). The shape and movement of fibroblasts in culture. In *Cell and Tissue Interactions*, edited by J. W. Lash and M. M. Burger. Raven Press, New York.
- Abercrombie, M., Flint, M. H., and James, D. W. (1956). Wound contraction in relation to collagen formation in scorbutic guinea-pigs. *Journal of Embryology and Experimental Morphology*, **4**, 167-175.
- Abercrombie, M., Heaysman, J. E. M., and Pegrum, S. M. (1971). The locomotion of fibroblasts in culture. IV. Electron microscopy of the leading lamella. *Experimental Cell Research*, **67**, 359-367.
- Bray, D., and Thomas, C. (1975). The actin content of fibroblasts. *Biochemical Journal*, **147**, 221-228.
- Carpenter, G., and Cohen, S. (1976). 125 I-labeled human epidermal growth factor: binding, internalization, and degradation in human fibroblasts. *Journal of Cell Biology*, **71**, 159-171.
- Conrad, G. W., and Hart, G. W. (1973). Do tissue-specific populations of fibroblasts arise during development? *Journal of Cell Biology*, **59**, abstract 63a.
- Fazzari, I. (1930). Sulla forma differente delle cellule mesenchimali dei vari organi nelle culture dei tessuti 'in vitro'. *Archiv für experimentelle Zellforschung besonders Gewebezuchtung*, **9**, 359-383.
- Franks, L. M., and Wilson, P. D. (1977). Origin and ultrastructure of cells *in vitro*. *International Review of Cytology*, **48**, 55-139.
- Gabbiani, G., and Montandon, D. (1977). Reparative processes in mammalian wound healing: the role of contractile phenomena. *International Review of Cytology*, **48**, 187-219.
- Gilula, N. B., Reeves, O. R., and Steinbach, A. (1972). Metabolic coupling, ionic coupling and cell contacts. *Nature (London)*, **235**, 262-265.
- Goldman, R. D., Lazarides, E., Pollack, R., and Weber, K. (1975). The distribution of actin on non-muscle cells. *Experimental Cell Research*, **90**, 333-344.
- Gospodarowicz, D., and Moran, J. S. (1975). Mitogenic effect of fibroblast growth factor on early passage cultures of human and murine fibroblasts. *Journal of Cell Biology*, **66**, 451-457.
- Green, H., Goldberg, B., and Todaro, G. J. (1966). Differentiated cell types and the regulation of collagen synthesis. *Nature (London)*, **212**, 631-633.
- Hayflick, L., and Moorhead, P. S. (1961). The serial cultivation of human diploid cell strains. *Experimental Cell Research*, **25**, 585-621.
- Heaysman, J. E. M., and Pegrum, S. M. (1973). Early contacts between fibroblasts. An ultrastructural study. *Experimental Cell Research*, **78**, 71-78.
- Holliday, R., Huscchtscha, L. I., Tarrant G. M., and Kirkwood, T. B. L. (1977). Testing the commitment theory of cellular aging. *Science*, **198**, 366-372.
- Isenberg, G., Rathke, P. C., Hülsmann, N., Franke, W. W., and Wohlfarth-Bottermann, K. E. (1976). Cytoplasmic actomyosin fibrils in tissue culture cells. Direct proof of contractility by visualisation of ATP-induced contraction in fibrils isolated by laser microbeam dissection. *Cell Tissue Research*, **166**, 427-443.
- Izzard, C. S., and Lochner, L. R. (1976). Cell-to-substrate contacts in living fibroblasts: an interference reflexion study with an evaluation of the technique. *Journal of Cell Science*, **21**, 129-160.
- Lewis, J. H., and Wolpert, L. (1976). The principle of non-equivalence in development. *Journal of Theoretical Biology*, **62**, 479-490.
- Manner, G. (1971). Cell division and collagen synthesis in cultured fibroblasts. *Experimental Cell Research*, **65**, 49-60.
- Millis, A. J. T., Hoyle, M., and Field, B. (1977). Human fibroblast conditioned media contains growth-promoting activities for low density cells. *Journal of Cellular Physiology*, **93**, 17-24.
- Papayannopoulou, T. G., and Martin, G. M. (1967). Alkaline phosphatase 'constitutive' clones: Evidence

- for *de novo* heterogeneity of established human skin fibroblast stains. *Experimental Cell Research*, **45**, 72-84.
- Pitts, J. D., and Sims, J. W. (1977). Permeability of junctions between animal cells. Intercellular transfer of nucleotides but not of macromolecules. *Experimental Cell Research*, **104**, 153-163.
- Rechler, M. M., Podskalny, J. M., and Nissley S. P. (1977). Characterization of the binding of multiplication-stimulating activity to a receptor for growth polypeptides in chick embryo fibroblasts. *Journal of Biological Chemistry*, **252**, 3898-3910.
- Rubenstein, P. A., and Spudich, J. A. (1977). Actin microheterogeneity in chick embryo fibroblasts. *Proceedings of the National Academy of Sciences of the United States of America*, **74**, 120-123.
- Schwarz, R., Colarusso, L., and Doty, P. (1976). Maintenance of differentiation in primary cultures of avian tendon cells. *Experimental Cell Research*, **102**, 63-71.
- Steinberg, J. (1973). The turnover of collagen in fibroblast cultures. *Journal of Cell Science*, **12**, 217-234.
- Stoker, M. G. P. (1973). Role of diffusion boundary layer in contact inhibition of growth. *Nature (London)*, **246**, 200-203.
- Stoker, M. G. P., O'Neill, C., Berryman, S., and Waxman V. (1968). Anchorage and growth regulation in normal and virus-transformed cells. *International Journal of Cancer*, **3**, 683-693.
- Summerbell, D., and Wolpert, L. (1972). Cell density and cell division in the early morphogenesis of the chick wing. *Nature New Biology*, **239**, 24-26.
- Wang, K., Ash, J. F., and Singer, S. J. (1975). Filamin, a new high-molecular-weight protein found in smooth muscle and non-muscle cells. *Proceedings of the National Academy of Sciences of the United States of America*, **72**, 4483-4486.
- Wessells, N. K. (1977). *Tissue Interactions and Development*. W. A. Benjamin, Menlo Park, California.
- Yamada, K. M., Yamada, S. S., and Pastan, I. (1977). Quantitation of a transformation-sensitive, adhesive cell surface glycoprotein. *Journal of Cell Biology*, **74**, 649-654.